

The background of the slide is a photograph of several surgeons in an operating room. They are wearing blue scrubs, white surgical masks, and blue bouffant caps. They are focused on a patient lying on the operating table, with their hands visible near the patient's head. The lighting is bright, typical of a surgical environment.

# Surgery miniOSCE

Done By: Yazan Alawneh

***Note: the answers in this file are based on the PYQ, so there is a possibility of inaccurate answers***

# Contents

Topic	Slide
Thoracic Surgery	3
Vascular Surgery	15
Genitourinary Tract	47
GI Tract (Esophagus, Stomach, Intestines)	52
Liver, Spleen, Pancreas, Gallbladder & The Adrenals	106
Anorectal Region	149
Bariatric Surgery	163
Salivary Glands	173
Neck & Thyroid	180
Breast	217
Pediatric Surgery	258
Skin	318
Burns	375
General Surgery & Others	391
Tools & Instruments	433

An anatomical illustration of the human thoracic cavity. The image shows a translucent blue human torso with the internal organs of the chest highlighted in red and pink. The trachea, bronchi, and lungs are clearly visible, along with the heart and major blood vessels. The rib cage and spine are also depicted. The word "Thoracic" is overlaid in large white letters with a black outline.

Thoracic

**Q: This is a chest X-Ray for a 35-years old female with a history of breast cancer 3 years ago, who presented to the clinic with progressive shortness of breath and cough.**

**Q1: What is the Dx?**

- Malignant Pleural Effusion

**Q2: What is the next step in Mx?**

- Tube thoracostomy (Chest tube)





## **Q1: What is the Dx?**

- Right sided hemothorax

## **Q2: Name 2 other findings?**

- 1) Absence of diaphragmatic angle
- 2) Right side multiple rib fractures
- 3) Right side clavicle fractures

## **Q3: What are the indication of needle thoracotomy after chest tube insertion?**

- initial loss  $>1.5$  L of blood
- Continuous blood loss of 200 ml per hour over 2-4 hour



**Q: Hx of motor vehicle accident (MVA):**

**Q1: What is the Dx?**

- Left sided hemothorax

**Q2: What is the Mx?**

- Chest tube insertion



**Q: A patient after a motor vehicle accident?**

**Q1: What is the Dx?**

- left sided hemothorax  
(obliterated costophrenic angle)

**Q2: What is the rapid initial Mx?**

- Needle decompression

**Q3: What is the definitive Mx?**

- Chest tube



**Q: A scuba diver came to ER, his CXR shows the following:**

**Q1: What is the immediate MX?**

- Needle thoracostomy

**Q2: Where to insert the needle?**

- 2<sup>nd</sup> intercostal space

**Q3: What is the procedure you want to do next?**

- Pleurodesis





### **Q1: What is the Dx?**

- Right sided tension pneumothorax

### **Q2: Mention 2 signs on CXR?**


- 1) Tracheal deviation
- 2) Left lung compressed or collapsed

### **Q3: Mention 2 signs on PE?**

- 1) Absent breath sounds in affected side
- 2) Jugular venous distention

### **Q4: What is the Mx?**

- Needle decompression
- Chest tube



**Q: 18 year old male presented with sudden progressive shortness of breath and underwent this investigation:**

**Q1: What is the Dx?**


- Spontaneous Pneumothorax


**Q2: What is the Mx?**

- Chest tube/needle

**Q3: Give 2 indications to do surgery?**

- Failure of decompression
- Hemo-pneumothorax





Tension Pneumothorax :  
The most reliable sign of tension pneumothorax is **depression of a hemidiaphragm.**




Pneumothorax in the Supine Patient . The **"deep sulcus sign"** is seen here (arrow) in the left lung base.

**right-sided pneumothorax  
with a chest tube inserted.**

- pneumothorax localizes more towards the apex of the lung.
- Notice that the markings are absent from the apex down to some degree.
- Notice the white visceral line.


Chest tube






# Surgical emphysema

- Radiolucent striations outlining pectoralis major
- It is usually benign, and treatment is directed at reversing the underlying cause.



It is unilateral  
**diaphragmatic  
paralysis.**(right)

we can still see the  
costodiaphragmatic  
angle so it is not  
effusion or  
hemothorax.



A detailed medical illustration showing a cross-section of a blood vessel. A catheter with a blue tip is inserted into the vessel. A stent, made of a mesh of grey rectangular cells, is positioned around the catheter. The vessel wall is shown in red and purple, with a yellowish, textured area representing plaque or a lesion. The word "Vascular" is written in large, white, bold letters across the center of the image.


Vascular

**What's A:** inferior epigastric artery

**What's B:** direct inguinal hernia

**What's C:** indirect inguinal hernia


**What's D:** femoral hernia





## Q: A Question was asking about the following arteries?

- 1- Left gastroepiploic artery
- 2- Gastroduodenal artery
- 3- Short gastric arteries



**Q: Patient had hip replacement 5 days ago:**

**Q1: What is the Dx?**

- DVT

**Q2: What is the Mx?**

- LMWH & Warfarin on discharge

**Q3: Mention 4 DDx?**

- 1) DVT
- 2) Cellulitis
- 3) Lymphadenopathy, lymphatic obstruction
- 4) Chronic Deep Vein Insufficiency
- 5) Rupture of baker's cyst

**Q4: What are the complications:**

- 1) Pulmonary embolism
- 2) Ulcers
- 3) Ischemia



**Q1: What is the Dx?**

- Varicose veins

**Q2: What is the system involved in this part (name the vessel)?**

- Great (long) Saphenous vein  
(Superficial Venous System)

**Q3: Name 2 modalities of Mx?**

- 1) High ligation and vein stripping
- 2) Sclerotherapy

**Q4: Mention 2 complications?**

- 1) Ulcers
- 2) Bleeding
- 3) Thrombophlebitis
- 4) Discomfort, pain



**Q5: Mention 2 minimally invasive procedures to do for this condition?**

- 1) Sclerotherapy
- 2) Radiofrequency Ablation
- 3) Endovenous Laser Ablation

**Q6: Best imaging test?**

- Doppler US or Venogram

**Q7: How to determine the level of defect in the varicose veins?**

- Turncate test





**Q1: What is this?**

- AV shunt

**Q2: Done in patients that undergoes what?**

- Hemodialysis

**Q3: What is the complication seen in the picture?**

- Aneurysm



**Q: A 60 year old female with CKD on hemodialysis:**

**Q1: What is the following complication?**

- Pseudoaneurysm

**Q2: Mention other complications that may occur?**

- Thrombosis, Steal syndrome, CHF, infection



**Q: Patient complained of abdominal pain and a pulsatile mass:**

**Q1: Name of this study?**

- Angiogram

**Q2: What is this pathology and where is its location?**

- AAA (Abdominal aortic aneurysm)  
near the bifurcation

**Q3: Mention 2 lines of Mx?**

- 1) Open surgical repair
- 2) Endovascular surgery






**Q1: Name of this study?**

- 3D angiography

**Q2: What is your Dx?**

- AAA



**Q: A patient with a hx of atrial fibrillation, presented with a sudden severe abdominal pain:**



**Q1: Name of this study?**

- CT Angiogram

**Q2: Dx?**

- AAA (Abdominal aortic aneurysm)

based on the Hx: Rupture AAA is more accurate





**Q1: What is the structure?**

- Abdominal Aorta

**Q2: What's the best repair method for this?**


- Stent





### Q3: What is the Mx (2 Mx modalities)?

- Medical or Surgical according to the size
  - 1) Endovascular repair
  - 2) Open repair



Abdominal x-ray with evidence of the **calcified edge** of the **abdominal aortic aneurysm**.



**Q: This is a CT Angio for the renal arteries:**

**Q1: What is the Dx?**

- Bilateral Renal Artery Stenosis

**Q2: What is your Mx?**

- Renal Angioplasty & Stenting



**Q: After RTA, the patient present with dilated veins?**

**Q1: Mention 2 causes?**

- 1) Pericardial Effusion
- 2) Cardiac Tamponade

**Q2: What is your Mx?**

- Pericardiocentesis



### Q1: What is the Dx?


Cardiac Tamponade

### Q2: What is the C/P that the patient come with?

- 1) Beck's triad :  
hypotension  
increased JVP  
muffled heart sounds.
- 2) Pericardial effusion
- 3) Kussmaul's sign.

### Q3: What is the Mx?

immediate decompression via  
needle pericardiocentesis.





**Q: Post-RTA patient came to ER, he was hypotensive with SOB:**

**Q1: What is the pathology?**


- Cardiac tamponade

**Q2: What is the next step in Mx?**

- Pericardiocentesis

**Q3: What is the consequence for this pathology?**

- Obstructive shock
- Pulmonary Edema
- Beck's Triad



**Q: a Pt experienced sudden severe pain radiating to the back:**

**Q1: What is the X-Ray finding?**

Widened Mediastinum

**Q2: What is the Dx?**


Aortic dissection

**Q3: What is the gold standard for Dx? And what is the disadvantage for it?**




Aortography, time consuming

**Q4: What is the Mx:**


- 1) Stanford A: Surgical
- 2) Stanford B: Medical (control BP)



## Classification of aortic dissection

			
Percentage	60%	10–15%	25–30%
Type	DeBakey I	DeBakey II	DeBakey III
	Stanford A (Proximal)		Stanford B (Distal)

**Westermarck's sign:** Decreased pulmonary vascular markings on CXR in a patient with pulmonary embolus



**Figure 1.** Chest radiograph demonstrating a prominent central pulmonary artery (early Fleishner's Sign, red arrows) and a cut-off of the pulmonary arteries bilaterally (Westermarck sign, black arrows).

# Mitral stenosis

## X-ray findings :


- Enlarged left atrium.
- **Straight line sign.**

## Diagnostic tests :

- Echocardiogram.
- Catheterization.

## Mx:

- Open heart surgery.
- Balloon valvoplasty.
- Valve replacement.



Mitral Stenosis



Source: <http://phil.cdc.gov>



## Q1: What does the arrow indicate?

Cervical rib

## Q2: What is your concern?

It can cause a form of thoracic outlet syndrome due to compression of the lower trunk of the brachial plexus or subclavian artery.

## Q3: What might the pt complain of?

- 1) parasthesias & numbness in the upper part mainly usually in 90% of cases are in the ulnar distribution.
- 2) Weakness manifested by difficulty grasping or holding a pen , this is a result of arterial and or neural compression.
- 3) The hand is usually cold.





**Q1: What is this sign?**

Raynaud's phenomenon.

**Q2: What is the most likely Dx?**

Buerger Disease

**Q1: What is the Dx?** Venous Ulcer

**Q2: What is the pathophysiology?**

- Blood stasis and increased Pressure inside the veins due to venous valves insufficiency

**Q3: if this happened after 5 days of surgery what is the main cause?**

DVT

**Q4: Risk of transformation to?** SCC

**Q5: Name 2 causes?**

- venous insufficiency and stasis (as DVT, varicose veins)

**Q6: What is the sign?**

- Lipodermatoseclerosis



**Q7: What is the most common site?**

Most Common site is lower 1/3 of the leg just above the medial malleolus.



**Q8: Name 2 points that goes with your Dx?**

- 1) Location: lower medial aspect of the leg
- 2) Hyperpigmentation around the ulcer





# Venous Ulcer Characteristics :

## where ?

\*Lower 1/3 of leg \*gaiter area \*anterior to medial malleolus.

## cause?

Commonly a history of:

\* (DVT) \*Obesity \*Calf muscle pump function deficits  
\*Valve incompetence in superficial perforating veins.

## description?

\*Ulcer has uneven edges \*Ruddy granulation tissue \*No dead tissue.  
\*Reddish brown pigmentation (Hemosiderin) \*Evidence of healed ulcers \*Edema that may leak and cause maceration, varicose eczema, itchy skin and scale  
\*Dilated and tortuous superficial veins \*Leg may be warm \*Hair on leg  
\*Normal leg and foot pulses.

## pain?

\*Moderate to no pain at all \*Pain if present is eased by raising the leg





**Q: A 75 year old male, heavy smoker, presented with this lesion.**

**Q1: Identify the lesion:**  
ischemic arterial ulcer

**Q2: Give two symptoms which might be associated with the condition:**

- 1) claudication
- 2) rest pain



# Arterial Leg Ulcer Characteristics

## where?

- \*At tips of toes or between toes
- \*Over phalangeal heads
- \*Above lateral malleolus, over the metatarsal heads, on the side or sole of feet.
- \* **MC distal end of the limbs**

## cause?

Commonly a history of:

- \*Aging
- \*Diabetes
- \*Arteriosclerosis
- \* Smoking
- \*Hypertension.

## description?

- \*Deep pale base
- \*Well defined edges
- \*Black or necrotic tissue
- \***Minimal / no hair**
- \***Thin, dry and shiny skin**
- \*Thickened toe nails
- \*Leg may be cool
- \*Leg becomes pale when elevated
- \*May have neuropathy
- \*Nil or diminished leg and foot pulses.
- \* **Punched out-appearance**

## Pain?

- \***Very Painful**
- \*Pain is reduced by lowering the leg to a dependent position.
- \* **Not palpable pulses**



**Q1: What is the most probable cause for this patient's condition?**

Lower Limb Ischemia

**Q2: What is the best imaging test to put a treatment plan?**

CT Angio, Angiogram,  
Doppler US



## Q1: What is the pathology?

- Gangrenous necrosis of the big toe

## Q2: Mention 4 signs of peripheral ischemic disease?

- 1) Pale
- 2) Hair loss
- 3) Cold
- 4) Pulselessness



**Remember the 6 P's of peripheral vascular disease:**

**P**allor

**P**ain

**P**aresthesia

**P**aralysis

**P**ulselessness

**P**oikilothermia

**Q: A patient walks 400 meters before feeling pain and having to rest, his job requires him to walk for 1 kilometer everyday, what do you do for this patient?**

- a) Lifestyle modification
- b) Medical therapy
- c) Bypass
- d) Angiogram **(correct answer)**



A medical professional in a white lab coat with a stethoscope around their neck is holding a tablet in their left hand and pointing with their right index finger at a digital illustration of the human genitourinary tract. The illustration shows the kidneys, ureters, bladder, and reproductive organs in a stylized, glowing red and orange color scheme with a wireframe overlay. The text "Genitourinary Tract" is superimposed over the illustration in a large, white, bold font with a black outline.


# Genitourinary Tract

## Q1: What is the imaging?


- MCUG

## Q2: Mx?


- Antibiotic for UTI
- Endoscopic injection
- Surgery




Grade I




Grade II



Grade III



Grade IV



Grade V

**Q1: What is the name of this study?**

- MCUG

**Q2: What is the name of this pathology (without abbreviation)?**

- Vesicouretral reflux (VUR)



**Q1: What is the pathology?**


- Left dilated tortuous ureter and hydronephrosis (right pic)

**Q2: What is the cause behind this?**

- Posterior urethral valve
- Congenital

**Q3: What are the 2 complications that might occur?**

- 1) Recurrent UTIs
- 2) Kidney scarring



**Q1: Name the finding?**

- Staghorn stone or Struvite stone

**Q2: What is the Etiology?**

- Urease producing bacteria (proteus, klebsiella, pseudomonas)






A person wearing a grey button-down shirt and a black belt is shown from the waist up. They are holding their right hand over their stomach, specifically the upper right quadrant, indicating abdominal pain. The background is a plain, light grey.

# Gastrointestinal Tract (Esophagus, Stomach & Intestines)

**Q: A 60 yo male patient  
came complaining of  
Dysphagia, halitosis,  
swelling in the neck:**

**Q1: What is the Dx?**  
Pharyngeal pouch

**Q2: How to Dx the pt?**  
Barium Swallow



**Q: Patient came complaining of dysphagia for both liquids & solids:**

**Q1: What is the sign?**

- Bird peak sign

**Q2: Name the study?**

- Barium swallow

**Q3: What is the definitive Dx?**


- Achalasia

**Q4: What is the definitive diagnostic test?**



- Manometry

**Q5: Mention 2 modalities of Mx?**

- 1) Esophageal sphincter (Hellers) Myotomy
- 2) Balloon dilation



May lead to esophageal carcinoma 2ry to Barrett's esophagus from food stasis.



**Q: a pt came complaining of dysphagia for both solids and liquids.**

**Q1: What is the Dx?**

Diffuse Esophageal Spasm (DES)

**Q2: What is the sign?**

corkscrew appearance

**Q3: How to Diagnose?**

- 1) Barium
- 2) Manometry (most accurate)

**Q4: What is the Mx?**

diltiazem or nifedipine and nitrates

**Q5: How to differentiate it from Nut-cracker esophagus?**

By manometry (the nut cracker: peristaltic contractions with high amplitude, while the DES is non-peristaltic with high contractions)





## **Q1: Define Barret's esophagus?**


Change in the normally squamous lining of the lower esophagus to columnar epithelium (metaplasia)

**Q2: What common type of cancer you will see?** Adenocarcinoma

**Q3: What is the cause?** Chronic GERD

**Q4: How to Dx?** Endoscopy

**Q5: Mx?** PPI and follow up





### **Q1: What is the Dx?**



Mallory Weiss Tear Syndrome

### **Q2: How to Diagnose it?**

Hx & Upper Endoscopy

### **Q3: Mx?**

It resolves spontaneously



**Q: Patient with Intermittent dysphagia for solids only with no pain:**

**Q1: What is the Dx?**

Schatzki ring (lower esophageal ring)

**Q2: Name an abnormality associated with it?**



Hiatal Hernia

**Q3: How to diagnose it?**

Barium swallow and endoscopy

**Q4: Mx?**

Dilation by bougie method or through the scope  
hydrostatic balloon, and the patients are placed  
on PPI after dilation



**Q: Patient with Intermittent dysphagia  
for solids only with no pain:**

**Q1: What is the Finding?**


Esophageal Webs  
(E.g. Plummer vinson syndrome)

**Q2: How to diagnose it?**

Barium swallow and endoscopy

**Q3: Mx?**

Dilation




# Plummer-Vinson syndrome:

1. Esophageal web
2. IDA
3. Dysphagia.
- 4. Spoon-shaped nails**
5. Atrophic oral & tongue mucosa.

\*especially occurs in elderly women; 10% develop squamous cell carcinoma.

\*May respond to treatment of IDA.





# Esophageal stricture

- Dysphagia : constant/ slowly progressive/ solids then liquids.
- Causes : long history of incomplete treated reflux/ prolonged NG tube placement/ lye ingestion.
  - Dx : barium swallow.
  - Treatment: dilation.





## Zenker's diverticulum:

- It is a **false diverticulum** (not involving all layers of the esophageal wall).
- Outpouching of the upper esophagus.
- **Halitosis** / food regurgitation/ dysphagia.
- Elderly.
- Dx : **barium swallow**/endoscopy and NG tube are contraindicated (risk of perforation).
- Treatment : **surgical resection**.




## Q1: What is the Dx?


Esophageal Varices

## Q2: Mx?


- 1) Therapeutic endoscopy
- 2) Ligation, banding,  
sclerotherapy
- 3)  $\beta$ -blockers (e.g. propranolol).



# Hiatal hernia



**Paraesophageal Hernia**



## Sliding hernia (type 1)

Mostly asymptomatic but can cause reflux

Complications : reflux > esophagitis > Barrett's esophagus > cancer/ aspiration pneumonia

Treatment: medical with antacids, PPI, H<sub>2</sub> blockers/ if failed : surgical (lap. Nissen fundoplication )

## Para esophageal hernia (2)

Dysphagia/ stasis gastric ulcer/ no reflux


Complications: hemorrhage/obstruction/ strangulation.

Treatment : surgical.

## Epiphrenic diverticulum


Presentation: Dysphagia to solid foods with upper abdominal discomfort.

Often associated with hiatal hernia.



# esophageal cancer

- is more after 50 years, most between 60-70 years.
- more in males.
- risk factors**: smoking, alcohol, and hot fluid drinkers.
- Relevant Hx**: GERD and Barrett's, stricture, Plummer Vinson syndrome, Celiac disease, Esophageal achalasia and diverticulum.
- common **symptoms** are dysphagia, reflux, weight loss, and mediastinal invasion symptoms (chest pain, hoarseness, etc.)
- they might also suffer from anemia due to nutritional deficiency.
- treatment** : surgical resection if small and localized.
- If large or Metastatic: combination of CTX and RTX prior to surgery.





easy mnemonic to remember esophageal CA risk factors  
ABCDEFGH:

A- Achalasia/Alcohol

B- Barrett's esophagus

C- Cigarettes

D- Diverticula

E- Esophageal web, stricture

F- Fat/Family hx

G- GERD

H- Hot liquid

# Gastric cancer


**Adenocarcinoma:** m.c type (95% ).

R.F: diet ( smoked meat , high nitrates , low fruits and vegetables) , smoking , family history , blood group type A , H. pylori , prev. partial gastrectomy , adenomatous gastric polyps , atrophic gastritis .


## Subtypes:

**diffuse type:** 70% ,from lamina propria, proximal , worse than intestinal type , invasive and Metz , in younger pt.

**intestinal type:** 30%,from gastric mucosa, distal , ass with H.pylori , well formed glandular structures.



Ulcerating adenocarcinoma




Intestinal type

**Diagnosis:** endoscopic with biopsy is the method of choice/ double contrast barium meal .

**Treatment:** surgical resection with wide margin >5cm and lymph nodes dissection .

If tumor is proximal or midbody do total gastrectomy with roux-en-y ,if tumor is distal do distal subtotal gastrectomy .



A. CT image of Linitis plastica (arrows denotes a thickened gastric wall).

**Linitis Plastica (leather bottle):**

when the entire stomach is involved and looks thickened .

**Q1: What is the Dx?**

Gastrointestinal Stromal Tumor  
(GIST)

**Q2: What is the MC site?**

Greater curvature (Stomach)

**Q3: What are the cells of origin?**

Cells of Cajal

**Q4: Name the Gene Mutation?**

C-KIT

**Q5: How to Mx?**


Resection

Chemo (Imatinib)

**Q6: How to Diagnose?**

CT / EGD/colonoscopy

Sx: abdominal pain/ nausea/  
abdominal mass/ occult GI bleeding.




**Q: A 50-years old male patient has recently become cachectic and developed ascites.**

**1. Name the findings on examination (lower picture) and CT scan (upper picture).**

- Sister Mary Joseph Nodule

**2. Mention 2 possible sources for this lesion.**

- GI cancers, Gynecological cancers, Melanoma





**Q: You are doing endoscopy and you found this lesion?**

**Q1: Describe what you see?**

- Comment on the shape, size, location, color, presence of necrosis, discharge, etc..

**Q2: What is the likely Dx?**

- Stomach cancer or ulcer

**Q3: Next step in Mx?**

- Biopsy



**Q: You are doing endoscopy and you found this lesion, pain is relieved by eating and exacerbated in empty stomach?**

**Q1: What is the likely Dx?**

- Peptic (duodenal) ulcer

**Q2: name 2 complications?**

- 1) Perforation
- 2) Bleeding



**Q1: What is A and B?**



**A** > Gastritis “not sure”

**B** > Duodenal Ulcer

**Q2: Name 2 causes?**

1) NSAID

2) H. Pylori



**Q: The patient presented with sudden severe pain and abdominal distension:**

**Q1: What is the sign?**

- Coffee bean sign



**Q2: Name the signs you?**

- 1) Dilated large bowel
- 2) Coffee bean sign

**Q3: What is your Dx?** Sigmoid volvulus

**Q4: What is the MC site?** in Sigmoid






### Q5: Mx?

- Resuscitation and untwist (detorsion) the bowel and go for surgery (this is done by means of sigmoidoscopy or colonoscopy)

### Q6: Mention 2 causes for this condition?

- Chronic constipation
- Sigmoid tumor





**Q1: What is the study?**


- Barium Enema

**Q2: What is the Dx?**

- Volvulus

**Q3: What is the Mx?**

- Detorsion




**Q1: What is the study?**

- Barium follow through

**Q2: What is the pathology?**

- Midgut volvulus




## Q1: What is the Dx?

- Volvulus (Midgut)

## Q2: If the bowel was viable and not gangrenous, what to do?

- Viable SB > Close and observe
- Other option: Ladd's Procedure



### **Q1: What is the study?**


- Barium follow through

### **Q2: What is the pathology?**

- Midgut volvulus due to malrotation

### **Q3: What is the Clinical ER Presentation?**

- acute abdominal pain , distention , constipation , vomiting



# Malrotation

normally the duodenojejunal junction is to the left of the spine. In malrotation it is to the right of the spine .





### Q1: What is the Dx?

Small intestinal obstruction

### Q2: What is the radiological findings?

Dilated bowel loops (Jejunal), and air in the rectum

### Q3: This is a picture of obstruction, Is it partial/complete? Why?

- Partial obstruction
- Because there is air in rectum

### Q4: What is the appearance?

Step-ladder appearance



**Q: A 30 year old female presented with sudden abdominal pain and fever and diffuse tenderness of the abdomen:**

**Q1: What is the Dx?**

Perforated viscus

**Q2: What is the radiological finding?**

Air under diaphragm

**Q3: What is the Mx?**


Laparotomy and exploration

**Q4: What is the mcc?**

Post-op

**Causes:**

1. Perforation of duodenal ulcer.
2. Following Laparoscopic procedure
3. Following Tubal Insufflation Test
4. Infection with gas forming organisms
5. Most common cause is post operative.
6. Chilaiditi's sign - due to interposition of colon between the Diaphragm and the Liver such a gas shadow can be obtained even in a normal individual.



**Q: A 55 years old patient with PUD came with forceful vomiting:**

**Q1: What is the pathology?**

- Gastric outlet obstruction (pyloric obstruction) – Pyloric Stenosis

**Q2: What is the electrolyte disturbances the patient has?**

- Hypokalemic hypochloremic metabolic alkalosis

**Q3: What is the gold standard for Dx?**



- US “not sure”

**Q4: Mention 2 causes?**

- 1) Gastric Carcinoma
- 2) Peptic ulcer disease (PUD)

**Q5: Name it's effect on ventilation?**

- Hypoventilation



**Q: A 48-years old patient presented with acute abdomen. PMH shows atrial fibrillation. Laparotomy was done:**

**Q1: What is the Dx?**


- Acute Mesenteric Ischemia

**Q2: What is the most affected artery in this condition?**

- Superior mesenteric artery

**Q3: Appropriate Mx?**

- Resection & Anastomosis





**Q1: What is the Dx?**

- Diverticulosis

**Q2: Mention 2 complications?**

- 1) Infection
- 2) Perforation
- 3) Obstruction

**Q3: What is the most common site?**

- Sigmoid






# Diverticulosis or Diverticular disease of the sigmoid colon


**Dx.** Colonoscopy

**Mx.** Mainly  
supportive: diet rich of  
fiber



# Colovesical fistula

- the most common cause is diverticulitis and it's the most common fistula formed in DD.
- other causes : colon CA , crohn's , radiotherapy ,trauma.
- This picture is double contrast barium enema.



diverticula

**Q: Female patient came complaining from fistulas and other symptoms and a colonoscopy was done:**

**Q1: What is the Dx?**

- Crohn's Disease

**Q2: What are the usual Sx?**

- Abdominal pain
- Fever with weight loss
- Diarrhea

**Q3: How do we treat those patients?**



- Azathioprine (6 mecaptopurine) + steroids





# Crohn's disease (IBD):

- Autoimmune disease
- SKIP LESIONS
- the m.csite is the terminal ileum,
- often no involvement of the rectum (in UC the rectum is always involved)
- Extraintestinal manifestations: arthritis , pyoderma gangrenosum ,erythema nodosum
- it involves the full thickness of the bowel wall , with the serosa ,mesentery and regional LNs ( while in UC it was only the mucosa that's involved)
- Macroscopically : the bowel wall is thick and red ( in UC it's very thin ), the mucosa has a cobblestone appearance
- Microscopically we will find non- caseating granulomas , with narrow deep fissure ulcers.
- Complications : strictures and fistulae ( in UC : hemorrhage , perforation , CA , and toxic megacolon)
- Radiology : Barium enema --> STRING SIGN
- Surgery plays a minor role in the treatment




**Q1: What is the Dx?**

- Ulcerative colitis

**Q2: Mention 2 drugs used in  
Mx?**

- 1) Steroid
- 2) Azathioprine





**Q: Known case of UC, with Hx of bloody diarrhea and abdominal pain:**

**Q1: What is the abnormality?**

- Transverse Toxic megacolon

**Q2: One complication?**

- Perforation
- Peritonitis



# Ulcerative colitis ( IBD )

UC is an autoimmune disease  
the rectum is always involved

\* smoking: protective.

- extracolonic manifestations :


arthritis ( sacroiliitis and ankylosing spondylitis), eyes ( iritis , keratitis ) , renal ( calculi & pyelonephritis , Skin (erythema nodosum & pyoderma gangrenosum), blood (anemia & higher risk of DVT), hepatic disease & cholangitis (PSC)

- investigations:

- if perforated --> Air under diaphragm on AXR
  - in chronic UC --> LEAD PIPE colon + and TOXIC MEGACOLON on AXR.

- Treatment :

- medical : mainly steroids ,/
  - Surgery (proctocolectomy with Brooke ileostomy ) is indicated when : medical treatment is failed , toxic megacolon , perforation and subsequent peritonitis , too frequent relapses , duration of more than 10 years ( >15 years --> 5% risk of CA )



Lead pipe colon



**Q1: What is the Dx?**

Colon Cancer

**Q2: What is the screening method?**

Colonoscopy


**Q3: What is the tumor marker?**

CEA


**Q4: What is the appearance?**

Apple-core

-Adenomatous polyps are precancerous.



### Apple-core Appearance of the Colon





# Gardner's Syndrome


( AD)

a familial adenomatous polyposis syndrome with cutaneous manifestations.

1) **Colonic polyps** ( hundreds with 100% risk of malignancy if untreated).


2) **Ostromas** ( the picture of an osteoma of the mandible).

3) **Lipomas and epidermoid cysts** ( on the forearm )






multiple small ulcers located  
in the distal duodenum in a  
patient with gastrinoma  
(**Zollinger- Ellison syndrome**)




# Q: What is the Dx?

## Gross Appendicitis



# Acute appendicitis

- Sx : pain (periumbilical area) >> nausea and vomiting >> anorexia >> pain migrates to RLQ (constant and intense, usually < 24 hrs.).
- Tenderness maximally at **McBurney's point**.
- Obturator sign/ psoas sign/ rovsingsign/ valentino sign.
- **Appendectomy** is the m.c.c of emergent abdominal surgery.
- Dx of ruptured appendix : fever >39 / high WBC/ rebound tenderness/ periappendiceal fluid collection on ultrasound.
- If normal appendix is found upon exploration, take it out ( even in chron's ).
- Appendiceal abscess : percutaneous drainage/antibiotics / elective surgery 6 wks later.



## **Q: Appendicitis Scenario:**

### **Q1: What is the pathology?**

- Acute Appendicitis

### **Q2: What is the name of it's scoring system?**

- Alvarado scoring system

### **Q3: What is the sequence of the pain?**

- Visceral somatic sequence of pain

### **Q4: Write 2 features found on US?**

- 1) Blind-ending tubular dilated structure >6mm
- 2) Appendicolith with acoustic shadow
- 3) Distinct appendiceal wall layers
- 4) Periappendiceal fluid collection
- 5) Periappendiceal reactive nodal enlargement




# Alvarado scoring system (Appendicitis)

Mnemonic (MANTRELS)	Value
Symptom	
Migration	1
Anorexia-acetone	1
Nausea-vomiting	1
Signs	
Tenderness in right lower quadrant	2
Rebound pain	1
Elevation of temperature $>37.3^{\circ}\text{C}$	1
Laboratory	
Leukocytosis	2
Shift to the left	1
Total score	10

## Q: What is the Dx?

- Peutz-Jeghers syndrome

- autosomal dominant.
- hereditary intestinal polyposis syndrome.
- hamartomatous polyps in the GI tract.
- **circumoral pigmented nevi.**





**Q1: What is your diagnosis ?**

FAP (focal adenomatous polyposis – in the colon & rectum)

**Q2: What is the cause of death before the age of 50?**

Cancer (untreated patients develop cancer by the age of 40-50)

**Q3: MOI?** Autosomal Dominant

**Q4: Associated tumors?** Duodenal Tumors

**Q5: Mx?** Total Proctocolectomy and ileostomy



**Q: patient with Hx of lower GI bleeding  
& this is the colonoscopy:**

**Q1: What is the Dx?**

- Angiodysplasia

**Q2: the Cause?**


- Degeneration of submucosal venous  
wall and formation of AVM

**Q3: the Mx?**

- 1) Laser
- 2) Electrocoagulation
- 3) Surgery

**Q4: What is the most common site?**

- the cecum or ascending colon





# Pseudomembranous colitis




Colonoscopy showing  
pseudomembranes

**cause:** *C. difficile*

**risk factors:** use of Antibiotics.

**diagnosis:** toxin assay in stool.

**treatment:** Metronidazole




- Abdominal CT.
- similarity between the thickened edematous wall of pseudomembranous colitis to that of an accordion.
- What is the sign?  
*Accordion sign.*

Name	Region & info	Indications
<b>Barium Swallow</b>	to visualize the area from the mouth to the stomach (esophagus)	a. Symptoms of gastro-esophageal reflux b. Dysphagia, related to: Esophageal (Web, stricture, tumor, achalasia), vascular abnormalities
<b>Barium Meal</b>	Double contrast (gas+barium) to visualize the stomach and the duodenum	a. Gastro-esophageal reflux b. Gastric or duodenal ulcer c. Hiatus hernia d. Gastric tumors
<b>Barium follow-through</b>	To visualize the small intestine, taken every 1/2 hr till we reach the large intestine (stool white)	a. IBS (crohns mostly) b. small bowel tumor/lymphoma (filling defect) c. Small bowel obstruction
<b>Barium Enema</b>	Double contrast (barium + air), to visualize the colon, and it's the only contrast given in the rectum (by Folly's)	a. Abdominal mass b. Large bowel obstruction / volvulus c. Diverticular disease d. Colonic tumor

An anatomical illustration of the human torso, focusing on the abdominal cavity. The rib cage is shown in a light beige color, with the ribs curving around the sides. The liver is a large, reddish-brown organ located in the upper right quadrant. The spleen is a smaller, reddish organ located in the upper left quadrant. The pancreas is a small, yellowish organ located in the center of the abdomen. The gallbladder is a small, pear-shaped organ located in the lower right quadrant. The adrenal glands are two small, yellowish organs located on top of the kidneys. The background is a solid light blue color.

# Liver, Spleen, Pancreas, Gallbladder & The Adrenals



**Q1: What is this triangle?**

- Calot's Triangle

**Q2: Name 3 borders?**


- 1) Inferior border of the liver
- 2) Cystic duct
- 3) Common hepatic duct



**Q: This 60-years old patient developed abdominal pain, bloody diarrhea and fever. He came back from a tour trip to a south west Asian country 3 weeks ago. CT was done.**

**1. What is the most likely diagnosis?** Liver Abscess (Ameobic)

**2. What is the treatment of choice?** Metronidazole



**Q: Name the following complications of liver cirrhosis:**

**A >** Ascites

**B >** Caput medusa (dilated veins)

**C >** Hematoma (easily bruised)




**Q1: What is the sign?** Caput Medusa

**Q2: What is the Dx?** Liver Cirrhosis



# Liver Abscess

- Pyogenic (bacterial “gram negative”) / parasitic (amebic) / fungal.
  - Most common site is right lobe.
- Treatment : pyogenic ( IV antibiotics + percutaneous drainage) / amebic (metronidazole+ drainage).
- Indications of surgical drainage in pyogenic : multiple lobulated abscesses/ multiple percutaneous attempts failed.
- Indications of surgical drainage in amebic: refractory to metronidazole/ bacterial co-infection/ peritoneal rupture.






**Q: Patient presented  
lethargic and febrile a week  
after a surgery for  
choolangitis:**

**Q1: What is your Dx?**

- Liver abscess

**Q2: Mx?**

- Percutaneous drainage, &
- Antibiotic administration



**Q: A 45 year old male presented with RUQ discomfort and pain, this is his abdominal CT.**

**Q1: What is the radiological finding?**

Peri-cyst and daughter cysts  
(hydatid cyst disease).




**Q2: Mention 2 complications:**

Rupture and anaphylaxis/  
obstructive jaundice.

**Q3: Give 2 drug that can be used?**

Albendazole, Mebendazole

is a **parasitic infestation** by a tapeworm of  
the genus **Echinococcus**.



**Q: Abdominal US image for a woman lives in rural area:**

**Q1: What is the name of this sign?**

- Water lily sign

**Q2: Most probable etiology for this sign?**

- Caused by tapeworm *Echinococcus granulosus*
- Another cause is *E. multilocularis*





# Caroli disease

is a congenital disorder comprising of multifocal cystic dilatation of segmental intrahepatic bile ducts.

presentation is in childhood or young adulthood. The simple type presents with RUQ pain and recurrent attacks of cholangitis with fever and jaundice.

Prognosis is generally poor. If disease is localized, segmentectomy or lobectomy may be offered. In diffuse disease management is generally with conservative measures; liver transplantation may be an option.






# Hepatic Hemangioma

- Most common benign solid tumor.
- Variants:
  - Capillary : m.c / <2cm /no need for surgery.
  - Cavernous : giant.
- Vague upper abdominal tenderness with no mass.
- **Not premalignant.**
- **Percutaneous biopsy is contraindicated** (risk of hemorrhage).
- U/S is the first test.
- MRI is the most sensitive & specific.



- Until recently, no medical therapy capable of reducing the size of hepatic hemangiomas had been described.
- Surgical treatment may be appropriate in cases of rapidly growing tumors. Surgery may also be warranted in cases where a hepatic hemangioma cannot be differentiated from hepatic malignancy on imaging studies.



# Hepatic Adenoma

## Risk factors:

Female/ birth control pills/ anabolic steroids/ glycogen storage disease.

it is estrogen sensitive  
(pregnancy may cause it to increase in size, OCP).




**Complications:** rupture with bleeding/ necrosis/ **risk of cancer.**

**Treatment:** if small, stop pills> it may regress> if not, surgical resection.  
If large or complicated : surgical resection



# Focal nodal hyperplasia


- Use of estrogen OCP may have a role.
- **Not premalignant.**
- Most are solitary, 20% multiple.
- Most common indication for surgery is inability to exclude malignancy.
- LFT : normal.
- Angiography : hypervascular mass with enlarged peripheral vessels and a single central feeding artery.
- **ttt : nucleation**/ diagnostic uncertainty will require an open excisional biopsy.



Classic CT finding: liver mass with central scar.


# Hepatocellular carcinoma (hepatoma)

- Most common 1ry malignant liver tumor.
- Risk factors: hepatitis B / cirrhosis/ Alfa toxin/ alpha 1 antitrypsin deficiency.
- Painful hepatomegaly.
- **Tumor marker: alpha fetoprotein.**
- Dx: needle biopsy with CT or U/S guidance.
- The m.c site of Metz :lungs.



**CT : black arrows (hepatoma)**





### Q1: What is the finding?

- Fluid in Morrison's pouch

### Q2: The Dx?

- Hemoperitoneum (blood)
- Ascitis (fluid)

*Morison's pouch:* The hepatorenal recess is the space that separates the liver from the right kidney.

**Q: a patient with RUQ pain:**

**Q1: What is the Dx?**


- Porcelain gallbladder

**Q2: What is the major risk?**

- Adenocarcinoma of  
gallbladder

**Q3: What is the Mx?**

- Elective Cholecystectomy



**Q: A 40 year old female patient after a bariatric surgery, presented with this US?**

**Q1: What is the Dx?**

- Gallstone

**Q2: What are the indications of performing a surgery in asymptomatic patient for this condition?**

- Porcelain gallbladder
- Congenital hemolytic anemia
- Gallstone >2.5 cm

**Q3: If the organ got inflamed where would be the pain and where it would radiate?**

- Pain would be in the RUQ, and radiate into the right subscapular area



# Gallbladder stones (Cholelithiasis)




Figure 1




Figure 2

Acoustic shadow

- 80% of patients are asymptomatic.
- Complications: acute and chronic cholecystitis/ CBD stones/ gallstone pancreatitis/ cholangitis.
- **U/S detects GB stones in more than 98% of cases.**
- Abdominal X-ray detects only 15%.
- If symptomatic/ complicated / asymptomatic but (sickle cell diseases, DM, pediatric, porcelain GB, immunosuppression) : cholecystectomy.





acoustic shadow

## Acute cholecystitis

- HIDA scan (the most accurate test).
- U/S (the diagnostic test of choice).
- Constant pain (not biliary colic).

### Sonographic findings in acute cholecystitis

- Impacted stone in cystic duct or GB neck
- Positive sonographic Murphy's sign
- Thickening of GB wall (**>3 mm**)
- Distention of GB lumen (**> 4 cm**)
- Pericholecystic fluid collections (frequent)
- Hyperemic GB wall on color Doppler (**supportive test**)

**None of above signs pathognomonic**

**Combination of multiple signs make correct diagnosis**

# Gallstone ileus


- occurs when a large gallbladder stone erodes into the duodenum via a fistula, eventually obstructing the ileal lumen usually some centimeters proximal to the ileocaecal junction.

On the X-ray :

1 radiopaque gallstone in the bowel.


2 gas in the gallbladder.

4- small bowel distention.



# emphysematous cholecystitis

- Gas forming bacteria (E.coli).
- Often results in perforation.
- Usually in males/ elderly/ DM.



**Q: After RTA, the patient present with left shoulder pain:**

**Q1: What is your Dx?**

- Splenic Rupture

**Q2: What is your Mx?**

- Splenectomy






Grade <sup>a</sup>	Type	Description of Injury
1	Hematoma	Subcapsular, < 10% surface area
	Laceration	Capsular tear, < 1 cm parenchymal depth
2	Hematoma	Subcapsular, 10–50% surface area Intraparenchymal, < 5 cm in diameter
	Laceration	1–3 cm parenchymal depth; does not involve a trabecular vessel
3	Hematoma	Subcapsular, > 50% surface area or expanding; ruptured subcapsular or parenchymal hematoma
	Laceration	> 3 cm parenchymal depth or involved trabecular vessels
4	Laceration	Laceration involving segmental or hilar vessels and producing major devascularization (> 25% of spleen)
5	Laceration	Completely shattered spleen
	Vascular	Hilar vascular injury that devascularizes spleen

Note—Adapted with permission from [2].

<sup>a</sup>Advance one grade for multiple injuries up to grade 3. The American Association for the Surgery of Trauma uses roman numerals.



**Q: RTA patient, HR = 130, he was hypotensive, a CT was done and shows the following?**

**Q1: How much blood did he loss?**

- Stage 3 hypovolemic shock – 30-40% - 1500-2000 ml

**Q2: What does the CT show?**

- Splenic Rupture




**Table 7-4 Signs and Symptoms of Advancing Stages of Hemorrhagic Shock**

	<b>Class I</b>	<b>Class II</b>	<b>Class III</b>	<b>Class IV</b>
Blood loss (mL)	Up to 750	750–1500	1500–2000	>2000
Blood loss (%BV)	Up to 15%	15–30%	30–40%	>40%
Pulse rate	<100	>100	>120	>140
Blood pressure	Normal	Normal	Decreased	Decreased
Pulse pressure (mmHg)	Normal or increased	Decreased	Decreased	Decreased
Respiratory rate	14–20	20–30	30–40	>35
Urine output (mL/h)	>30	20–30	5–15	Negligible
CNS/mental status	Slightly anxious	Mildly anxious	Anxious and confused	Confused and lethargic

BV = blood volume; CNS = central nervous system.

# Acute Pancreatitis

- Cut off sign and Ileus.
- White arrow points to Transverse colon cut off at Splenic flexure.
- No air in descending colon.
- TC: Transverse colon.
- I: Represents small bowel loops with air suggestive of Ileus.



Causes : gallstones/ ethanol/ trauma/ steroids/ mumps/ autoimmune/ scorpion bite/ hyperlipidemia/ drugs (diuretics, INH)/ ERCP.

Treatment : supportive (90% resolve spontaneously)



**Q: A 45-years old male patient, alcoholic, presented with a 24-hour history of upper abdominal pain and repeated vomiting. On examination of the abdomen, he was found to have these signs.**

**Q1: Name those signs?**

**A > Cullen's**



**B > Grey Turner's**

**Q2: Mention 2 causes?**

- Any retroperitoneal hemorrhage

1) Acute hemorrhagic pancreatitis

2) Abdominal trauma bleeding from aortic rupture



# Chronic Pancreatitis

**most common cause is chronic alcoholism.**

Abdomen x-ray showing pancreatic calcifications.




# Pancreatic necrosis

- Dx: abdominal CT with contrast.
- Dead pancreatic tissue doesn't take up the contrast.



# Pancreatic pseudocyst

- **The m.c.c is chronic alcoholic pancreatitis.**
- findings : high amylase/ fluid filled mass on ultrasound/
- it is a collection of fluid rich in pancreatic enzymes, blood, and necrotic tissue.
- to exclude malignancy >>you have to check the level of CA 19-9 ( tumor marker).
- Complications: bleeding into the cyst/ infection/ pancreatic ascites.



- **If not resolved spontaneously within 6 weeks : drainage.**



**Q1: What is the type of imaging?**

- MRCP

**Q2: Mention 2 abnormalities?**


- 1) Stone in the CBD  
(arrow – filling defect)
- 2) Dilated CBD



**Q1: What is the study?** MRCP

**Q2: The structure pointed?** Pancreatic duct (Stricture)

**Q3: What is the next step?** ERCP



**Q: 60 year old female with RUQ pain and fever.**

**Q1: Identify this type of image:**  
MRCP

**Q2: Give two radiological findings:**  
CBD stone shadow/ CBD dilation.

**Q3: What is your diagnosis?**  
Ascending cholangitis.



# Choledocolithiasis

- Common bile duct stones.
- ERCP (the diagnostic test of choice, also therapeutic).
- If ERCP fails, CBD is opened surgically and stones removed.

The huge tube is the endoscope. It is going down from the esophagus, through the stomach, to the duodenum (1st then 2nd parts), and stops near the ampulla of Vater.

A tube in the endoscope is pushed into the ampulla and fills the CBD with a dye. X-ray is taken.

As you can see, there is a black shadow stone in the CBD.





**Q1: What is the name of this investigation?** ERCP

**Q2: Mention two abnormalities seen in this picture:**

Filling defect & distended common bile duct



**Q1: What is the type of imaging?**

- ERCP

**Q2: Indications?**

- Obstructive jaundice

**Q3: Complications of ERCP?**

- Pancreatitis

**Q4: Mention 2 findings?**

- 1) Dilated CBD
- 2) Multiple stones



### **Q1: What is the Dx?**

- Primary sclerosing cholangitis  
(Beading)

### **Q2: Which disease is associated with it?**


- Ulcerative colitis

### **Q3: Which type of malignancy the patient may develop?**

- Cholangiocarcinoma

### **Q4: Diagnostic test?**

- ERCP



**Q: a patient with thyroid medullary cancer, & a CT was done:**

**Q1: What is your next step? (not sure what the dr. meant so here is the possibilities):**

- Assess the functionality of the adrenal tumor by hx, physical ex and ordering lab tests: KFT (Na, K, Creatinine, Urea) / Aldosterone levels/ cortisol/ metanephrine / noremetanephrine / vanillyl mandelic acid (VMA)
- pheochromocytoma
- 24h urine analysis for catecholamine metabolites (VMA/Meta)

**Q2: If the patient has no genetic abnormality and the lesion is not functioning what will you do next?**

- Because it is very large > surgery adrenalectomy, the dr said : If it was more than 4 cm then you have to remove it immediately





**Q: a patient presented with episodic sweating and hypertension:**

**Q1: What is the Dx?**

- Pheochromocytoma

**Q2: What is the 1<sup>st</sup> thing to do?**

- Check if functional or not by checking cortisol, renin, angiotensin and VMA,... etc

**Q3: What raise the possibility of malignancy?**

- >4 cm
- necrosis
- hemorrhage

**Q2: What is the size that would be considered an indication for surgery?**

- >4 cm



**Q: Lab investigations show high aldosterone level and high ratio of PAC to PRA:**

**Q1: What is your Dx?**

- Conn's tumor

**Q2: Mention a common presentation for this patient?**

- Hypertension



Functional adrenal tumors can cause several problems depending on the hormone released. These problems include:

### 1. Cushing's Syndrome:

This condition occurs when the tumor leads to excessive secretion of cortisol. While most cases of Cushing's Syndrome are caused by tumors in the pituitary gland in the brain, some happen because of adrenal tumors. **Symptoms of this disorder include diabetes, high blood pressure, obesity and sexual dysfunction.**

### 2. Conn's Disease:

This condition occurs when the tumor leads to excessive secretion of aldosterone. **Symptoms include personality changes, excessive urination, high blood pressure, constipation and weakness.**

### 3. Pheochromocytoma:

This condition occurs when the tumor leads to excessive secretion of adrenaline and noradrenaline. **Symptoms include sweating, high blood pressure, headache, anxiety, weakness and weight loss.**

**Q: A 40-years-old female,  
previously healthy, presented  
with acute abdominal pain,  
fever and itching**

**1. What is the diagnosis?**

Ascending Cholangitis

**2. What is the next imaging test  
to order for this patient?**

MRCP, ERCP

**3. Why is she having itching?**

Bile salts accumulation






# Pneumobilia

( Air in the biliary tree )

## Causes :

- Recent biliary instrumentation (e.g. ERCP or PTC)
- Incompetent sphincter of Oddi (e.g. sphincterotomy, following passage of gallstone.)
- Biliary-enteric surgical anastomosis.
- Spontaneous biliary-enteric fistula (cholecystoduodenal accounts for ~70% ).
- Infection (rare) (e.g. ascending cholangitis, anaerobes).



A person wearing a red hoodie and blue jeans is sitting on a blue surface. They are holding their right knee with both hands, suggesting discomfort or pain. The word "Anorectal" is overlaid in large white text with a black outline.


**Anorectal**

**Q: About the anatomy of anal canal:**

**A:** External anal sphincter

**B:** Internal anal sphincter

**C:** Dentate line



**Q: Patient has anal pain and itching:**


**Q1: What type of anal condition in this area (Area A)?**

- Ischiorectal abscess

**Q2: What is the Mx?**

- Cruciate incision with drainage with drainage of pus (without antibiotic)

- Extra: we use antibiotic in: systemic inflammatory response or sepsis extensive cellulitis, diabetes, immunosuppression







**1st Degree: No Prolapse**  
Just prominent vessels




**2nd Degree: Prolapse (come out) with strain**  
but spontaneously reduce (go back in)




**3rd Degree: Prolapse with strain**  
and have to be pushed back in




**4th Degree: Prolapsed out and**  
cannot be reduced or pushed back in







- a: superficial fistula
- b: intersphincteric fistula
- c: transsphincteric fistula
- d: suprasphincteric fistula
- e: extrasphincteric fistula



**Q: This is a 35-years-old patient c/o severe anal area pain**

**1. What is the diagnosis?** Perianal Abscess

**2. What is the treatment?** Drainage & Antibiotics Cover

**3. What is the possible sequel for this condition?** Fistula





**Q: A 25 year old male presented with anal pain and fresh blood PR, the peri-anal area is shown:**

**Q1: What is the Dx?** Bleeding Hemorrhoids

**Q2: What do you recommend?**

1) Bath sitz 2) Laxatives 3) High-fiber diet

**Q3: Beside bleeding, name 2 more complications?**

1) thrombosis 2) Infection 3) Ulcers

**Classification:** Internal (above dentate line)  
external (below dentate line).

**Risk factors:** constipation/ straining/  
pregnancy/ ascites/ portal HTN.

**Hemorrhoidectomy:**

- \* contraindicated in chron's.
- \* complications: pelvic infection/ anal stricture/ incontinence.



## Q1: Name the Dx?


- Pilonidal Sinus (PNS)

## Q2: Name 4 sites for it?

- 1) Inter-digital space
- 2) Natal cleft
- 3) Between breast
- 4) Axilla

**Treatment** If your PNS does get infected, surgery will most likely be recommended and may include the following:

- 1) Incision and Drainage
- 2) Wide Excision (reduce your chances of a reinfection. However; Your wound may take a long time to heal)
- 3) Excision and Primary Closure (reinfection chances are higher)



**Q: A 22-years old male patient presented with upper natal cleft area increasing in pain for the last 3 days.**

**1. What is your diagnosis?**

Gluteal Cleft Abscess of a Pilonidal Sinus

**2. What is the treatment?**

Incision & Drainage



# Fistula –in- ano

- From rectum to anal skin.
- Causes:  
anal crypt infection  
perianal abscess.
- Sx :  
perianal drainage  
itching  
diaper rash.





**Q: This pt has painful defecation:**


**1. Name the findings on examination of the anal area.**

A > Anal Fissure

B > Sentinel Pile

**2. Mention 2 treatment options.**


- **Lifestyle modification** with high fiber diet and increase fluid intake
- **Medical Management** (Laxatives, stool softeners, local anesthetic creams, botulinum toxin injection, sitzbath...etc)
- **Surgical Management** (Sphincter dilatation, Lateral internal sphincterotomy, Fissurectomy)




This is a chronic fissure with hypertrophic papilla & pile formation, the guidelines state that for chronic fissures medical management with botulinum toxin, stool softeners and anesthetic creams is indicated first. If the fissure is refractory to medical management then surgical intervention with lateral internal sphincterotomy is highly indicated, but sphincter dilatation could also be used.

# Anal fissure

- **Hypertonic internal sphincter.**
- Chron's disease may cause it.
- Very painful.
- Posterior fissures more common than anterior ones.
- Signs : **sentinel tag/ hypertrophied papilla/** blood on toilet paper.
- Surgery indication: chronic fissure / refractory to conservative treatment.
- Surgery: lateral internal sphincterectomy.
- Triad of chronic fissure: sentinel pile/ hypertrophied papilla/hypertonic sphincter.



Hypertrophic papilla



Sentinel pile

# Perianal warts

- Cause : condylomata acuminata (HPV).
- The major risk is SCC.
- Treatment : if small, topical **podophyllin**/ if large, surgical resection or laser ablation.








# Bariatric Surgery



- Weight reduction surgery for the morbidly obese.
- Morbid obesity : BMI > 40 or BMI > 35 with a medical problem related to morbid obesity (sleep apnea/ CAD/ DM/ HTN/ pulmonary disease/ breast cancer/ colon cancer/ arthritis/ sex hormone abnormalities/ venous stasis ulcers).



**Q1: Name this surgery?**

- Gastric bypass


(Single Anastomosis Gastric Bypass)

**Q2: Mention 2 types?**


- 1) Gastrojejunostomy
- 2) Duadenoileostomy

**Q3: What BMI is an indication for a surgery in a DM patient?**

- >35



# Lap Sleeve Gastrectomies (LSG)



**Q: A Patient that needed to reduce weight ASAP, and this surgery was done:**

**Q1: Which procedure is this?**

- Gastric Sleeve

**Q2: 2 Complications for it?**

- 1) Blood clots.
- 2) Gallstones
- 3) Hernia.
- 4) Internal bleeding
- 5) Leakage.
- 6) Perforation
- 7) Stricture







## Q1: Name this surgery?


- Roux-en-y gastric bypass (RYGB)

## Q2: Mention 2 mechanisms?

- 1) Malabsorption  
(Decrease gastric absorption)
- 2) Less space for food  
(early satiety)




# Roux-en-Y gastric bypasses (RYGB)



Gastric Balloon




# Gastric Band (LABG)





# Mini Gastric Bypass

## Gastric Bypass



An anatomical illustration of a human head in profile, facing right. The skin is semi-transparent, revealing the underlying structures. The parotid gland is shown as a large, pink, lobulated mass in the lower part of the face. The sublingual gland is visible as a smaller, similar mass above the lower lip. The submandibular gland is located below the jawline. A network of yellow nerves is depicted branching out from the glands and extending towards the eye and ear. The background is a solid dark grey.

# Salivary Glands

**Q1: What is the organ affected?**

- Parotid gland

**Q2: What is the most likely Dx?**

- Parotid Pleomorphic Adenoma

**Q3: What is the most common subtype?**

- Myxoid (not sure)

**Q4: What is 1 sign that will confirm your Dx?**

- Rubbery-hard, does not fluctuate and of limited mobility on physical examination

- **Benign** salivary gland tumor.
- The most common salivary gland tumor.
- Usual location : parotid gland.
- single firm, mobile, well- circumscribed mass.
- **Painless.**
- Slow growing.






## Q5: How do we treat this pt?

- Superficial parotidectomy, some said total parotidectomy

## Q6: Histology?

Epithelial  
Myoepithelial  
Stroma  
Pseudopods  
No true capsule





**Q: a patient had a superficial parotidectomy:**


**Q1: What is the most likely indication?**

- Parotid gland tumor (most likely pleomorphic adenoma)

**Q2: What is the nerve in risk of being damaged?**

- Facial nerve

Some said: great auricular nerve




**Q: 50 yo pt presented with bilateral neck swelling:**

**Q1: What is the Dx?**


- Warthin's tumor

**Q2: What is the malignancy risk?**

- 0.3%



- is the 2<sup>nd</sup> mc benign salivary gland tumor.
- More in males.
- Associated with smoking.
- **Only in parotid.**
- Usually at parotid tail.
- Cystic mass.



**Q1: if a surgery was done  
what is the nerve at risk to  
be injured?**

- Marginal Mandibular Nerve

**Q2: What is the risk of  
malignancy?**

-50%



Salivary Gland	Malignancy Rate	Incidence of Tumor
Parotid	20%	80%
Submandibular	50%	15%
Sublingual & Minor	70%	5%






# Sialolithiasis = salivary stones

## Submandibular salivary gland stone

- The stone is located in the Wharton's duct (most common site) :  
in the floor of the mouth near the frenulum of the tongue.







# Neck & Thyroid

# DDx of neck lumps

	Midline	Lateral
<b>Neoplastic</b>	Thyroid Parathyroid Pharyngeal/Laryngeal	Most tumors (lymphoma, carotid...)
<b>Congenital</b>	Thyroglossal duct cyst Laryngocele	Cystic Hygroma Branchial cleft cyst
<b>Infectious</b>	Ludwig's Angina	Most infections (cat-scratch, mononucleosis, sialadenitis...)
<b>Inflammatory</b>	Submental reactive lymphadenopathy Thyroiditis	Most reactive lymphadenopathy

## **Q1: What is the Dx?**

- Lacerated neck wound

## **Q2: What zone?**

- Zone 2

## **Q3: Name the borders for it?**

- From the angle of the mandible to the cricoid cartilage

## **Q4: When to intubate the patient?**

- 1) Expanding hematoma
- 2) Obstructive complication
- 3) Cervical vertebrae injury



## PENETRATING NECK INJURIES

---

What depth of neck injury must be further evaluated?

Penetrating injury through the platysma

Define the anatomy of the neck by trauma zones:

Zone III


Angle of the mandible and up

Zone II

Angle of the mandible to the cricoid cartilage

Zone I

Below the cricoid cartilage





How do most surgeons treat penetrating neck injuries (those that penetrate the platysma) by neck zone:

Zone III

Selective exploration

Zone II

Surgical exploration vs. selective exploration

Zone I

Selective exploration

What is selective exploration?

Selective exploration is based on diagnostic studies that include A-gram or CT A-gram, bronchoscopy, esophagoscopy

What are the indications for surgical exploration in all penetrating neck wounds (Zones I, II, III)?

**"Hard signs"** of significant neck damage: **shock**, exsanguinating hemorrhage, expanding hematoma, pulsatile hematoma, neurologic injury, subQ

## Bethesda diagnostic category

**VERY COMMON QUESTION!**

## Risk of malignancy

## Usual management

<b>I</b>	<b>Nondiagnostic or unsatisfactory</b>	Cyst fluid only Virtually acellular specimen Other (obscuring blood, clotting artifact, etc.)	1% to 4%	Repeat FNA with ultrasound guidance
<b>II</b>	<b>Benign</b>	Consistent with a benign follicular nodule (includes adenomatoid nodule, colloid nodule, etc.) Consistent with lymphocytic (Hashimoto) thyroiditis in the proper clinical context Consistent with granulomatous (subacute) thyroiditis Other	0% to 3%	Clinical follow-up
<b>III</b>	<b>Atypia of undetermined significance or follicular lesion of undetermined significance</b>		5% to 15%	Repeat FNA
<b>IV</b>	<b>Follicular neoplasm or suspicious for a follicular neoplasm</b>	Specify if Hurthle cell (oncocytic) type	15% to 30%	Surgical lobectomy
<b>V</b>	<b>Suspicious for malignancy</b>	Suspicious for papillary carcinoma Suspicious for medullary carcinoma Suspicious for metastatic carcinoma Suspicious for lymphoma Other	60% to 75%	Near-total thyroidectomy or surgical lobectomy
<b>VI</b>	<b>Malignant</b>	Papillary thyroid carcinoma Poorly differentiated carcinoma Medullary thyroid carcinoma Undifferentiated (anaplastic) carcinoma Squamous cell carcinoma Carcinoma with mixed features (specify) Metastatic carcinoma Non-Hodgkin lymphoma Other	97% to 99%	Near-total thyroidectomy

### Q1: What is the Dx?

- Thyroglossal duct cyst



### Q2: What is the structure on U/S (involved bone)?

- Hyoid bone

### Q3: What is the Mx?

- Sistrunk's procedure

(if the hyoid bone not removed the recurrence rate is > 50-60%)



**Q4: What is the malignancy risk?**

- 2%

**Q5: Name the malignancy that does not occur here?**

- Medullary Ca

**Q6: Complications?**

- Infection, malignant risk

**Q7: Sign to confirm your Dx?**

- Movement with tongue protrusion

**Q8: What is the risk of recurrence?**

- Sistrunk procedure reduces the recurrence risk from 60% to < 10%






**Q: This is the US of a 20 yo male with a neck lump.**

**1. What is the next step in approaching his condition? FNAC**


**2. What is the most likely Dx? Thyroglossal Duct Cyst**



**Q: This patient underwent surgery for the pathology depicted by the yellow arrow. Histology reported a malignancy of non-thyroid origin.**

**What is the most likely malignancy? SCC**


**What structure does the red arrow point to? Hyoid bone**



**Q1: Name the triangle of the neck in which the lesion is situated:**

anterior triangle.

**Q2: Give 2 DDx for the lump:**  
sialodenitis/ lipoma.




# Ludwig angina

pus accumulation in the **submental triangle**. causes pressure on the larynx and epiglottis and suffocation.

treated surgically by opening the submental area and draining the pus.





LUDWIGS  
ANGINA






# Carotid body tumor : in carotid triangle

- moves side by side.
- Dx: carotid angiogram.
- Surgical excision and preoperative embolization.
  - Lateral mass.





## Branchial cyst

- Smooth surface and globular.
- At the level of junction between upper and middle 1/3 of SCM.



## Branchial fistula

- formed by the 2<sup>nd</sup> branchial cleft and pouch.
- lined by ciliated columnar epithelium.
- Discharge : mucus or muco-pus.
- in anterior triangle.
- at junction between middle and lower third of SCM.
- congenital.
- surgery (excision).



## Sublingual dermoid cyst

- Medline congenital mass.
- Contents : hair follicles/ sebaceous cyst/ sweat glands.

## Plunging ranula



**Ranula** : cystic mucosa extravasation from sublingual salivary gland.

**Plunging** : if extended through mylohyoid muscle.

Treatment : excision.

**Q: Hx that suggest a thyroid nodule:**

**Q1: What is the Dx?**

- Multi-nodular goiter

**Q2: How to approach the patient with this Dx?**

- TFT
- US





## **Q1: What is the Dx?**

- Graves disease

## **Q2: Mention 2 signs that you can see?**

- Exophthalmos
- Significant hair loss
- Lid retraction



## **Q3: What is the 1<sup>st</sup> Sx patient will develop if she develops ophthalmoplagia?**

- Diplopia or Proptosis (not sure)

## **Q4: What is a drug you can give this patient before getting into surgery?**

- PTU (Propyl thiouracil), propranolol



**Q: 50 year old female patient  
present with hypothermia:**

**Q1: What is the endocrine  
disorder?**

- Hypothyroidism


**Q2: Mention 3 signs on face?**

- 1) Puffy face
- 2) Periorbital edema
- 3) Coarse hair





**A. Normal**



**B. Graves' disease**



**C. Toxic mng**



**D. Toxic adenoma**

**Q: Patient with hyper diffuse functioning thyroid:**

**Q1: What is the Dx?**

- Graves Disease

**Q2: What is the serological marker?**

- TSI (thyroid stimulating immunoglobulin)


**Q3: Mention 3 lines of Mx?**

1) Anti-thyroid drugs (carbimazole) +  $\beta$ -blockers

2) Radio-iodine

3) Surgery

\*\* All 3 are considered 1<sup>st</sup> line Mx





**Q1: What is the pathology?**


- Papillary Thyroid Carcinoma



**Q2: What is the rate of the malignancy?**

- 97-99%


**Q3: Mention 2 features seen in the picture?**

- 1) Nuclear Crowding
- 2) Orphan Annie Nuclei





- Papillary thyroid carcinoma:**
- a. Nuclear groove (blue arrow).**
  - b. Psammoma body.**



**Papillary thyroid carcinoma:**  
**(Intranuclear cytoplasmic inclusions)**



**Q1: What type of thyroid cancer do you expect to see in this patient?**

- Medullary

**Q2: What's the marker?**

- Calcitonin






**Q1: What type of thyroid cancer do you expect to see in this patient?**

- Medullary cancer

**Q2: Before surgery what type you must exclude?**

- MEN 2 (Pheochromocytoma)




**Q: Hx of thyroid nodule, US showing: micro-calcifications, investigation of blood vessels and reactive LN:**

**Q1: Bethesda Grade?**

- Bethesda 6

**Q2: What is your Mx?**



- Total Thyroidectomy



**Q: Images A & B demonstrate thyroid nodules that are considered sonographically suspicious for malignancy. Name the feature labelling each nodule suspicious.**

**A > Heterogeneous**



**B > Calcification**



**Q: What shall you do in the following cases ?**

**A. Thyroid** → repeat cytology

**B. Parathyroid** → removal (parathyroid adenoma)






## **Q1: Name the study?**


- Sestamibi scan of parathyroid

## **Q2: What is the most common cause of the condition?**

- Adenoma



**15 minutes**




**2 hours**

## Q1: Name the study?

- Sestamibi scan

## Q2: What is the pathology you see?

- Hyperfunctioning parathyroid glands



**Q1: Risk of disease to be from single nodule?**


- 85-90% Adenoma

**Q2: What is your Dx?**

- Single parathyroid gland adenoma

**Q3: What is your Mx?**

- Removal




## Q1: What is the Dx?

- Parathyroid adenoma (1ry hyperparathyroidism)

## Q2: The 1<sup>st</sup> Sx to develop if the patient had high PTH & Calcium?

- Bone pain (Since it's Hyper)
- if Hypo: Peri-oral numbness, carpal spasm






**Q: A 60-years old female complains of pain in her bones. She presents with a palpable central neck lump below the cricoid cartilage that moves upward upon swallowing.**

**1. What does the lump mostly represent?**

Parathyroid Carcinoma

**2. What is the bone condition called?**

Osteitis Fibrosia Cystica



**Q1: Name the Dx?**


- Parathyroid hot nodule

**Q2: Name the Rx?**

- Surgery (Lobectomy)

**Q3: Risk of malignancy?**


- Low risk (<3-5%)



**Q: Hx of palpable neck mass, recurrent renal stone, high level of calcium and parathyroid hormone:**

**Q1: Name the Dx?**

- Parathyroid carcinoma



**Q2: What is the minimal Mx to be done?**

- Parathyroidectomy or en-bloc resection of the parathyroid mass and any adjacent tissues that have been invaded by tumor . (from uptodate)

\*\*\* Note: En-bloc resection could include the ipsilateral thyroid lobe, paratracheal alveolar and lymphatic tissue, the thymus or some of the neck muscles, and in some instances, the recurrent laryngeal nerve

**Q: The morning post-total thyroidectomy the patient developed the sign seen in this figure:**

**Q1: Name of the sign?**

- Trousseau Sign

**Q2: What is the cause?**


- Hypocalcemia after removal of parathyroid glands

**Q3: What is the most likely cause of hypoparathyroidism?**

- Ischemic Injury



**Trousseau's sign** : Carpal spasm after occlusion of blood to the forearm with a BP cuff in pt with **hypocalcaemia**.







**Q1: What are the signs?**

- Chvostek and Trousseau signs

**Q2: What is the cation that influx and cause this sign?**

- Na<sup>+</sup> Sodium



# Breast

<https://radiologyassistant.nl/breast/bi-rads-for-mammography-and-ultrasound-2013>

## BI-RADS CATEGORIES

**BI-RADS 0 (incomplete):** Recommend additional imaging -- mammogram or targeted ultrasound

**BI-RADS 1 (negative):** Routine breast MR screening if cumulative lifetime risk  $\geq 20\%$

**BI-RADS 2 (benign):** Routine breast MR screening if cumulative lifetime risk  $\geq 20\%$

**BI-RADS 3 (probably benign):** Short-interval (6-month) follow-up

**BI-RADS 4 (suspicious):** Tissue diagnosis

**BI-RADS 5 (highly suggestive of malignancy):** Tissue diagnosis

**BI-RADS 6 (known biopsy-proven malignancy):** Surgical excision when clinically appropriate

## Final Assessment Categories

Category		Management	Likelihood of cancer
0	Need additional imaging or prior examinations	Recall for additional imaging and/or await prior examinations	n/a
1	Negative	Routine screening	Essentially 0%
2	Benign	Routine screening	Essentially 0%
3	Probably Benign	Short interval-follow-up (6 month) or continued	>0 % but ≤ 2%
4	Suspicious	Tissue diagnosis	4a. low suspicion for malignancy (>2% to ≤ 10%) 4b. moderate suspicion for malignancy (>10% to ≤ 50%) 4c. high suspicion for malignancy (>50% to <95%)
5	Highly suggestive of malignancy	Tissue diagnosis	≥95%
6	Known biopsy-proven	Surgical excision when clinical appropriate	n/a



## **FNAC (Breast)**

**C1: Unsatisfactory**

**C2: Benign**


**C3: Atypical cells**

**C4: Suspicious cells**

**C5: Malignant**

## ACR classification of breast density

ACR = American College of Radiology



There are four categories of mammographic density :

**ACR 1** : almost entirely fatty.

**ACR 2** : scattered areas of fibroglandular density.

**ACR 3** : heterogeneously dense.

**ACR 4** : extremely dense.

Metrics	Results	ACR type	Density percentage value (%)	Sensitivity (%)	Specificity (%)	Accuracy (%)
TP	97	1 (fatty breast)	<10	90.65	73.59	85.00
FP	14					
TN	39					
FN	10					
TP	66	2 (Fibro-glandular dense)	25-50	61.68	90.57	71.25
FP	5					
TN	48					
FN	41					
TP	22	3 (Heterogeneous dense)	50-75	20.56	96.23	45.63
FP	2					
TN	51					
FN	85					
TP	6	4 (Extremely dense)	75>	5.61	98.11	36.25
FP	1					
TN	52					
FN	101					

## TNM Class

## Criteria

T0 No evidence of primary tumor

T1a Carcinoma in situ

T1 < or = 2 cm

T1m1c microinvasion .1 cm or less

T1a >.1 to .5 cm

T1b >.5 to 1 cm

T1c >1 to 2 cm

T2 >2 to 5 cm

T3 >5cm

T4 Any size tumor with direct extension to : a) Chest wall or b) skin

T4a Chest wall, not including pectoralis muscle

T4b Skin edema, ulceration, satellite skin nodule

T4c 4a and 4b

T4d Inflammatory carcinoma



TNM Class	Criteria
Nx	Regional lymph nodes cannot be removed
N0	No regional lymph node metastasis
N1	<input type="checkbox"/> Metastasis to movable ipsilateral axillary lymph nodes <input type="checkbox"/> 1–3 ALN
N2	<input type="checkbox"/> Metastases in ipsilateral axillary lymph nodes fixed or matted (N2a) or met. only in clinically apparent ipsilateral mammary nodes without clinically evident axillary lymph nodes. ( N2b) <input type="checkbox"/> 4–9 ALN
N3	<input type="checkbox"/> Metastases in ipsilateral axillary or infraclavicular lymph nodes (N3a) or clinically apparent ipsilateral internal mammary lymph nodes (N3b) or ipsilateral supraclavicular lymph nodes (N3c) <input type="checkbox"/> 10 or more ALN
MX	Distant metastasis cannot be assessed
M0	No distant metastasis
M1	Distant metastasis

**Q1: What is the finding?** Male breast nipple changes

**Q2: Most common gene mutation associated with male breast cancer?** BRCA 2



**Q: A nipple biopsy for a female patient shows large cells with a clear cytoplasm, high grade nuclei and prominent nucleoli:**

**Q1: What is your Dx?**

- Paget disease of the breast/nipple (PDB)

**Q2: Mention 2 immuno-histochemical tests to differentiate it from melanoma?**

- 1) CEA (pos. in PDB)
- 2) Protein S100 (neg. in PDB)



### **Q1: What is the Dx?**

- Breast mastitis, Abscess

### **Q2: MCC?**

- S. Aureus

### **Q3: Mx?**

- Abx
- Incision & Drainage





**Q: 50 yo female has breast pain,  
breast only shows skin redness?**

**Q1: What is the Dx?**

- Inflammatory breast cancer

**Q2: Diagnostic procedure?**

- Tissue biopsy

**Q3: Mx?**

- Mastectomy + Radiotherapy

**Q4: What is the modality of Dx?**

- Triple assessment
- Mammogram + US

**Q5: According to TNM stage system  
the T stage is?**

- T4d






**Nipple retraction**  
(inversion).



**Peau d' orange**  
(orange peel).



**Skin dimpling**




**Paget disease** of the nipple  
(eczema around the nipple)

# Duct ectasia

- AKA Plasma cell mastitis.
- Condition Mimics cancer (nipple retraction, inversion, pain, Nipple discharge).
- disorder of peri- or post-menopausal age.
- Self-limiting condition.



duct ectasia :bilateral inversion and displaying transverse slit pattern






# Fine needle aspiration (FNA)

## \*\* Advantages :

- done in office
- minimal discomfort.

## \*\* Disadvantage :


- may not always rule out cancer when it's negative.






# Incisional biopsy

- Local anesthesia, often with mild sedation.
- Only part of the tumor is removed for Dx.
- Outpatient procedure.
- Done when the tumor is large.




# Excisional biopsy

- **The mc biopsy procedure.**
- Outpatient procedure.
- The entire lump is taken out using a small incision.



# Lumpectomy

- Excisional biopsy may be sufficient for the lumpectomy, if the margins were negative.
- With radiation therapy, it is as effective as modified radical mastectomy.





# Radiotherapy

## Side effects (self limited)

skin reddening & irritation/ darkening of the skin/ blistering/ minimal ↓ in blood counts/ mild fatigue/ lymphedema in the arm ( arm sleeves are used to control the swelling).



# Chemotherapy

## Side effects

hair loss/ ↓ blood counts/ nausea & vomiting/ ↓ platelet count when high dose is used/ mouth sores/ diarrhea/ loss of appetite/ wt gain/ menopause.



**Q1: What is the pathology?**

- Carcinoma en cuirasse

**Q2: What is its TMN?**

- Stage 4




**Q: Name the following views for mammogram:**

- Craniocaudal (CC)
- Mediolateral Oblique (MLO)

Craniocaudal (CC)

Mediolateral oblique  
(MLO)





**Q1: Name the study?**

- Mammogram

**Q2: Mention 2 abnormalities?**


- Mass with irregular border and  
calcification

**Q3: What is the Dx?**

- Breast Ca

**Q4: How to confirm your Dx?**

- Biopsy



**Q1: What is this view?**

- Mediolateral oblique

**Q2: What is this structure (arrow)?**


- Pectoralis major muscle

**Q3: What are the malignant changes seen on mammograms? Mention 3?**

1) Calcifications

2) Speculations

3) Mass with greater density than normal tissue



**Q: A 23-year-old single female presented to the clinic with rapidly growing (9cm) left breast mass over the last 6 months. The mass was irregular, hard and fixed at the time of examination:**

**Q1: Your Dx?**

- Phyllodes tumor

**Q2: What is this structure (arrow)?**

- Pectoralis major muscle

**Q3: if it is malignant, what is the common route of METS?**

- Hematogenous

**Q4: The mc site of METS?**

- Lungs



**Q: Female with ACR of 4 and BIRAD 0:**

**Q1: What is the % of breast density?**

- >75%

**Q2: What to do next?**

- Birads score: requires further investigations




## **Q: Breast with Birad 2:**

**Q1: What is the next step in Mx?**

- Routine screening

**Q2: What is the view in B?**


- Mediolateral oblique view



**Q: A 37-year-old female presented with right breast pain for the last 3 months. A breast ultrasound showed these findings consistent with BIRAD 4c.**


**Q1: The likelihood of malignancy is: 50-90%**

**Q2: The clinical T stage “if a diagnosis of invasive carcinoma is proved” is: T4**



**Q: A 40-years old married female presented with a right breast mass for 1-year duration. The patient had a history of a right breast mass excision 3 years ago. Physical exam showed a 4cm hard right breast mass which is fixed to the chest wall & the skin. Mammogram and ultrasound were consistent with BIRADS 5.**

- 1. Based on the TNM, the clinical T stage for this patient is? T4c**
- 2. The likelihood of malignancy based on imaging findings is? >95%**



**\*\*T4a : to chest wall only\*\***

**\*\*T4b : to skin only\*\***

**\*\*T4c : to both\*\***

**\*\*T4d: Inflammatory breast cancer\*\***

**Q1: What is the pathology?**

- Infiltrative ductal carcinoma

**Q2: What is its TMN?**

- Stage T3

**Q3: What is the sign?**

- Peau'd orange and nipple retraction, skin dimpling

**Q4: Give 2 DDx?**

- 1) Invasive ductal carcinoma
- 2) Inflammatory breast cancer



**Q5: What is the cause of this?**

- Invasion of lymphatics, causing lymph nodes obstruction



**Q: A pt came complaining of a tender cord like subcutaneous structure, pain, swelling and redness of the left breast:**

**Q1: Dx?** Mondor's Disease (Superficial Thrombophlebitis)

**Q2: What is the Mx?**

- NSAIDS
- Usually benign and self-limiting condition



**Q1: What is the name of this study?**

- Mammogram

**Q2: Mention 2 signs you see.**

- 1) Speculated mass
- 2) Microcalcifications

**Q3: What is the Dx?**

- Infiltrative Ductal Carcinoma



## **Q1: What is the pathology?**


- Phyllodes tumor (Brodie's)

## **Q2: What is the Mx?**

- Wide local excision

## **Q3: What is the like hood (%) of this tumor to be benign?**

- 90% benign



**Q: Female with mobile, mouse like lump in one breast:**

**Q1: What is the Dx?**

- Fibroadenoma

**Q2: What is the stage according to FNA?**

- C2



C1 = unsatisfactory.

C2 = cells present all benign; no suspicious features.

C3 = cells suspicious but probably benign.

C4 = cells suspicious but probably malignant.

C5 = Definitely malignant.



**Q: a 35 yo female patient:**

**Q1: What is the Dx?**

- Breast Cyst

**Q2: Name the sign (black arrow)?**

- Acoustic enhancement



**Q3: What are the indications for a biopsy in this female?**

1) Bloody aspiration

2) Failure to completely resolve

3) Recurrence after 2<sup>nd</sup> aspiration

4) Atypical cells



### **Q1: Describe the discharge?**

- Uniductal Bloody Discharge

### **Q2: What is the pathology?**

- Intraductal papilloma

### **Q3: Give a DDx?**

- Intraductal papilloma
  - Duct Ectasia
- Ductal invasive carcinoma

### **Q3: 2 imaging studies?**

- 1) Ductogram, Ductoscope
- 2) Mammogram, US

### **Q4: What is the risk of malignancy of this lesion?**

- 15%




**Q1: What is the mechanism  
that the breast cancer causes  
hypercalcemia?**


- Parathyroid hormone - related  
protein  
(not due to osteoclastic METS)

**\*\* Note:** The main pathogenesis of hypercalcemia in malignancy is increased osteoclastic bone resorption, which can occur with or without bone metastases. The enhanced bone resorption is mainly secondary to PTH-related protein






## Breast Cyst



**breast cancer:**  
dense mass with a  
spiculated margin.




**clustered microcalcification:** five or more calcifications, each measuring less 1mm in one cubic cm, the possibility of malignancy increases as a size of individual calcification decreases and the total number of calcification per limit area increases.



## The 2 major signs of malignancy in mammography:

1. Mass with spiculated margins or stellate appearance ( the single arrow ).
2. Microcalcifications (the double arrows ).




# Breast Infiltrating ductal cancer ultrasound.



This shows an irregular ductal  
tumor with nodules  
infiltrating the area around it.

# Sentinel Lymph Node



The sentinel nodes are the first place that [cancer](#) is likely to spread.




**Q1: What are the skin changes indicative of breast cancer in this image?**

Nipple retraction

Peau d'orange

**Q2: What is this procedure?**

Core needle biopsy  
(true-cut biopsy)



# Lymphangiosarcoma

- As a complication of long- standing lymphedema , usually in the edematous arm of post radical mastectomy patient.
- to prevent it : use elastic compression stockings.





# Pediatric Surgery

**Q: This 1 year old baby had this lesion since birth:**

**Q1: What is the most likely Dx?**  
Hemangioma

**Q2: What is the best Mx?**  
Observation and reassurance





# Vascular malformation



## Sturge weber syndrome

port wine stain vascular malformation involving the ophthalmic division.

- Usually not evident at birth.

### mnemonic :

S : seizures / U: unilateral weakness  
R: retardation ( mental ) / G: Glaucoma  
E : other eye problems



**Capillary hemangioma** in the eyelid obstructing the eye , might lead to Amblyopia "lazy eye".



The same patient at different ages (hemangioma)

hemangioma	Vascular malformation
Start as small lesions at the age of 3-4 months	seen at birth but may appear late
Grow to reach their maximum size at the age of 1 year then involution	Grow parallel to the child's growth
Female to male (3:1)	Female to male (1:1)
Rarely to cause any complications	High flow can lead to destructive changes
Spontaneous resolution unless complicated you should treat	Treatment : surgery/laser/ embolization

# Bilateral cleft lip and palate

## Cleft lip:

No functional deformity, only cosmetic deformity and **surgery is done at age of 3 months**.  
Breast feeding is not contraindicated.

## Cleft palate:

baby can't feed, can't speak and may lose his hearing by time (acquired).  
**surgery is done at age of 1 year** as a compromise between not losing his speaking abilities and the normal growth of face.



Unilateral incomplete



Unilateral complete



Bilateral complete



Incomplete cleft palate



Unilateral complete lip and palate



Bilateral complete lip and palate





# Pentalogy of Cantrell

1. Omphalocele.
2. Anterior diaphragmatic hernia.
3. Sternal cleft.
4. Ectopia cordis.
5. Intracardiac defect.



**Q1: What is the Dx?** Prune belly syndrome

**Q2: Mention 2 associated anomalies?**

- 1) Undescended testicles
- 2) Urinary tract abnormality such as unusually large ureters, distended bladder, Vesicoureteral reflux, frequent UTI's
- 3) VSD
- 4) Malrotation of the gut
- 5) Club foot

- thin flaccid abdominal wall.
- AKA eagle Barrett syndrome.
- absent abdominal wall musculature.
- dilation of bladder, ureter and renal collecting system.
- **95% in Males.**



# Bickwith-Wiedman syndrome

1. Macrosomia.
2. Macroglossia.
3. Organomegaly.
4. Abdominal wall defects.
5. Embryonal tumors.





# Torticollis

- Tilted neck.
- Causes:
  - 1) congenital** ( due to abnormal position of the fetus in uterus which leads to fibrosis of sternocleidomastoid muscle >> shortness of this muscle)

2) **acquired** : due to trauma leads to muscle spasm on one side/ fibrosis of SCM due to any cause.

3) **infection: lymphadenitis**



- Occurs at any age but most common in the 1<sup>st</sup> few months of life.
- Palpable hard mass in 1/3 of patients.
- The baby usually sleeps on the same side >> craniofacial deformity.
- Treatment : conservative using physiotherapy for 2-3 months.
- If no improvement, surgery is indicated (SCM myotomy).




# Cystic hygroma

- Fluid-filled sacs caused by blockages in the lymphatic system.
- **most hygromas appear by age 2.**
- **soft, non-tender, compressible lump.**
- high recurrence rate.
- usually located in the posterior triangle of the neck.
- **transillumination.**
- DDx: teratoma/hemangioma/
- encephalocele.




# Congenital malformations

**Think of Albort Syndrome**



# Esophageal atresia and tracheoesophageal fistula



# Manifestations of esophageal atresia:

- 1) **Upper part:** drooling of saliva/ bubbling of the saliva/ respiratory distress/ choking/ failure to pass nasogastric tube.
- 2) **Lower part:** accumulation of secretions which will lead to regurgitation and vomiting/ ischemia>> physiological death>> biological death (necrosis) >> rupture.

\* The more distal the obstruction, the more the distention of the lumen and so the more the possibility of rupture.




Neonates with esophageal atresia usually develop copious, fine white frothy bubbles of mucus in the mouth and nose. Secretions recur despite suctioning.



# Esophageal atresia and tracheoesophageal fistula

- Atresia of the upper esophagus evidenced by failure to pass a feeding tube.
- Gas in the abdomen.
- These findings are likely due to a esophageal atresia with a distal tracheoesophageal fistula (Type C TEF).




## **Q: New born x-ray, cyanosis and distressed:**

### **Q1: What is your Dx?**

- Tracheoesophageal fistula (because of the cyanosis)

### **Q2: Characteristic sign?**

- Failure to pass the nasogastric tube



**Q: A new-born baby had inability to swallow milk and frothy mouth secretions, this is his x-ray.**

**Q1: Mention two radiological signs?**  
inability to pass nasogastric tube/air in the stomach.

**Q2: What is the diagnosis?**  
Esophageal atresia with tracheo-esophageal fistula.



# ARDS

(bilateral diffuse  
pulmonary infiltrates )

## **Other DDx:**

- 1-severe pulmonary edema.
- 2-pulmonary hemorrhage.
- 3-pulmonary fibrosis.

( history differentiates  
between these conditions)

**Ground Glass Appearance**







This chest X-ray shows **air trapping** indicating **foreign body aspiration**.


It is the most common radiological sign shown on the X-ray after F.B aspiration.

Whenever you suspect F.B aspiration you have to do **bronchoscopy**.





**Tetralogy of Fallot**  
"boot" shaped heart on  
chest X-ray.



**Transposition of great  
vessels**  
Egg shaped heart

# Congenital diaphragmatic hernia

- X-ray of the abdomen and chest.
- features :
  - scaphoid abdomen.
  - **bowel is located in the left side of the chest.**
  - mediastinal shift towards the right.
- mortality is mostly due to pulmonary hypoplasia.
- Diagnosis: In prenatal period (ultrasonography)



## • Types :

### 1) Bochdalek hernia


(mostly on left side): posterolateral, mc.

### 2) Morgagni hernia

(mostly on the right side): retrosternal.

Hiatus hernia.

# Neonate with a prenatally diagnosed left congenital diaphragmatic hernia pre surgery.





	Omphalocele	Gastroschisis
Incidence	1:6,000-10,000	1:20,000-30,000
Delivery	Vaginal or CS	CS
Covering Sac	Present	Absent
Size of Defect	Small or large	Small
Cord Location	Onto the sac	On abdominal wall
Bowel	Normal	Edematous, matted

	Omphalocele	Gastroschisis
Other Organs	Liver often out	Rare
Prematurity	10-20%	50-60%
IUGR	Less common	Common
NEC	If sac is ruptured	18%
Associated Anomalies	>50%	10-15%
Treatment	Often primary	Often staged
Prognosis	20%-70%	70-90%

**Q1: What is the Dx?** Gastroschisis

**Q2: Name the procedure?** Silo

**Q3: The prognosis depends on?**

- Bowel status

**Q4: The indication of this procedure?**

- if the bowel is inflamed and primary closure is not possible
- to prevent dehydration, hypothermia, contamination

- **location : lateral to the umbilicus ( to the right ).**
  - defect size : 2-4 cm.
  - no sac.
  - cord is normally inserted into umbilicus.
- **contents : only bowel (edematous and matted ).**
  - GIT function : **prolonged ileus.**
  - associated anomalies : infrequent.





# Q1: What is the Dx?

- Omphalocele

# Q2: How is the GI function?

- Normal



- location : umbilical ring.
- The protrusion is covered by peritoneum.
  - defect size : >10 cm.
- cord : inserted into the sac.
  - GIT function is normal.
- contents : bowel +/- liver.
  - malrotation : present.
- associated anomalies : common (30-70 % ).





**Q1: What is the diagnosis in A,B?**


**A > Omphalocele**

**B > Gastroschisis**



**Q2: Which of these are more associated with congenital anomalies?**

**- Omphalocele**



### Q3: What is the 1<sup>st</sup> aid Mx for both?

- Carefully wrap in saline-soaked pads.
- Support without tension.
  - NG tube.
- Abdominal ultrasound.



**Q: Malrotation:**



**Q1: What's A and B?**

**A > Non-Rotation**

**B > Incomplete Rotation**

**Q2: Which one is the most commonly associated with volvulus?**

**- B**



**Q: What is the Dx according to:**

**A: Preterm baby > Necrotizing enterocolitis (NEC)**

**B: Full-term baby > Hirschsprung disease**





# Intussusception

- It is a cause of intestinal obstruction.
- M : F ( 3:2)
- In a previously healthy infant.
- (5 months - 3 yrs) idiopathic / (>3yrs) 2ry.
- m.c.c of I.O in the age of (5 months-3 yrs)
- Sudden onset, abdominal colic, vomiting.
- begins proximal to ileo-cecal junction.
- **Ba enema ( diagnostic and therapeutic).**
- The part that prolapses into the other is called the intussusceptum, and the part that receives it is called the intussusciptient.



**Q1: What is the investigation?**

- Abdominal US

**Q2: Name of the sign?**

- Target sign

**Q3: What is the pathology?**




- Intussusception

**Q4: How do we treat those patients in uncomplicated cases (stable)?/1<sup>st</sup> line of Mx?**

- Resuscitation, Hydrostatic (pressure) reduction using gas air or barium enema







Red currant jelly


Stool

RUQ Target sign (doughnut sign).

Pseudokidney sign






# • INTUSSUSCEPTION •



# Pyloric stenosis

- M : F (4:1)
- Age (3-6 wks)
- Progressive, persistent, projectile, non-bilious vomiting.
- Succation splash.
- Olive sign (enlarged pylorus is palpable).
- Hypochloremic alkalosis.
- Dx by abdominal U/S
- Higher risk when mother is affected.
- Surgical ttt: Ramstad's pyloromyotomy.
- No recurrence after surgery.





### **Q1: What is this?**

- Meckel's Diverticulum

### **Q2: Name 2 complications?**

- 1) Intestinal hemorrhage
- 2) Intestinal obstruction
- 3) Diverticulitis

### **Q3: Mention one common ectopic tissue you can find?**

- Gastric and pancreatic tissues




# Q4: Is it a true or pseudo-diverticulum?

## - True Congenital Diverticulum

-A memory aid is the rule of 2s:  
2% (of the population).  
2 feet (proximal to the ileocecal valve).  
2 inches (in length).  
2 types of common ectopic tissue (gastric and pancreatic)  
2 years is the most common age at clinical presentation  
2:1 male: female ratio



# Types of intestinal atresia



**Q1: What is the Dx?**

Jejunal atresia.

**Q2: Age of presentation?**

Neonate (till one month)


**Q3: How would u manage?**

Admit to NIC

fluid resuscitation

Antibiotic

NG suction and parental nutrition.




**Q: Intra-op image of a baby with symptoms of obstruction.**

**Q1: Give two findings:**

Dilated proximal loop,  
collapsed distal loop.

**Q2: What is the diagnosis?**

Type 1 intestinal atresia.






- **Apple peel intestinal atresia**  
(also type IIIb or **Christmas tree atresia**).

- Due to vascular accident.

- All the intestine is atretic, and forms a loop around the superior mesenteric artery.



# Intestinal obstruction




- Abdominal X-ray.
- Double bubble sign.
- represents dilation of the proximal duodenum & stomach.
- DDX : duodenal stenosis (mostly in the 2<sup>nd</sup> part of duodenum) / duodenal atresia.



Multiple air fluid levels seen in mechanical intestinal obstruction.

# Meconium ileus

- Intestinal obstruction from solid meconium concretions.
- >95% have cystic fibrosis.
- Sx: bilious vomiting/ abdominal distention/ failure to pass meconium.



# Hirschsprung's disease

- Congenital megacolon.
- It is an absence of ganglion cells distal in the bowel.
- Contracted non-peristaltic affected segment and a dilated hypertrophied proximal segment.
- M:F (4:1)
- Failure to pass meconium in the 1<sup>st</sup> 24-48 hrs of life.
- When compared to habitual constipation (no soiling/ no anal fissures).
- DDx : hypothyroidism/ sepsis.



Plain abdominal X-ray : dilated loops of bowel/ air-fluid level.



Barium enema study: **funnel shaped appearance** of colon ( megacolon – transitional zone- the affected narrowed segment).



**Q: A neonate failed to pass meconium, so a barium enema was done and shows this:**

**Q1: What is the Dx?**

- Hirschsprung disease

**Q2: What does the arrow indicate?**

- Transition zone

**Q3: What is the diagnostic test?**

- Biopsy
- Full thickness or rectal suction


**Q4: Name the radiology study?**

- Barium enema





# Umbilical Hernia

- more common in blacks.
- familial tendency.
- repair is carried out if closure does not occur by the end of 2<sup>nd</sup> year of life.
- repair performed after the age of 2 and before the age of 10.
- associated anomalies :
  - hypothyroidism.
  - hurler syndrome.
  - beckwith-wiedman syndrome.



# Patent urachus

- It is a remnant presents as **fistula** connecting the umbilicus & urinary bladder.
- Patients with prune belly syndrome have a patent urachus.
- Other forms : blind sinus/ cyst/ abscess.



(A) Prolapsed bladder was shown through the patent urachus. (B) Catheterization through the urethral orifice confirmed the communication between patent urachus and the bladder (black arrow: catheter tip).

# Vesicointestinal fissure

The terminal ileum is herniating through the cecum forming the so called **elephant trunk deformity**.



**Fig.1:** Showing omphalocele, lateral bladder plate, caecal plate and prolapsed ileum.



# Omphalitis

- Inflammation of the umbilicus.
- Occurs only in newborns.
- Can be fatal because of portal vein thrombosis.
- Infection can spread to the abdominal wall.
- Antibiotics and intensive care.



# Bladder Extrophy

- Defective enfolding of caudal folds.
- Associated with prolapsed vagina or rectum / **epispadias** / bifid clitoris or penis.




# Vesicoureteral reflux

- **Presentation : either antenatal hydronephrosis or clinical UTI.**
- Diagnosis : urine culture/ ultrasound/ **voiding cystourethrogram.**
- Nuclear cystogram for screening.
- DMSA scan to detect kidney scarring.
- Urodynamic study for lower urinary tract abnormalities (neurogenic bladder).



Spot film taken during VCUG shows unilateral grade 4 vesicoureteral reflux

# UVR grades



## Treatment :

- Spontaneous resolution is common in young children (only antibiotics).
- Indications for surgery: grade 4 and 5/ poor compliance with medications/ breakthrough febrile UTI despite adequate antibiotic prophylaxis/ poor renal growth/ kidney scars/ mild or moderate reflux in females that persist during puberty despite several yrs of observation.



### **Q1: What is the pathology?**

- Right scrotal swelling (Hemi-scrotal swelling)

### **Q2: Give two benign DDx?**

- Inguinal hernia, hydrocele

### **Q3: What is the name of peritoneal part remain patent?**

- Patent processus vaginalis



# Inguinal hernia

- Due to patent processus vaginalis.
- More common at the right side.
- Bilateral hernias occur in 5-15% of children with hernia.
- **Uncomplicated hernia** will bulge when the baby cry and reduces when the baby is relaxed , sleeping. Etc.
- Uncomplicated hernia must be operated (herniotomy).
- Herniotomy must be performed ASAP.
- 10-15% of children with on the other side. hernia on one side will develop a hernia



- **Complicated hernia** presents in the ER with pain/ management : resuscitation, reduce hernia, then repair within 24-48 hrs. ( as we fear strangulation and testicular atrophy).

## Q1: What is the Dx?

- Epispadias and Hypospadias


## Q2: Mention 2 associated anomalies?

- 1) Bladder extrophy
- 2) Bifid penis
- 3) Rectum prolapse


## Q3: Name 2 commonly associated features with this pathology other than the abnormally located urethral meatus:

- 1) Chordee  
(downward bending of the penis)
- 2) Hooded appearance of the penis

**Hypospadias**



**Epispadias**



## Q1: What is the Dx?

- Hypospadias

## Q2: What is the classification?

- 1) Anterior (50%)
- 2) Bifid Middle (30%)
- 3) Posterior (20%)

## Q3: When is the surgery performed?

6 – 18 months of age





- Glanular (opening on the glans) is the most common.



**Epispadias:** urethral opening is on the dorsal surface with abnormal penis. It is usually a part of a syndrome includes extrophy of the urinary bladder.

- Extremely rare.





Copyright the Lucina Foundation, all rights reserved.

**Q: This is a 5 yo boy.**

**Q1: Give two clinical findings:**

scrotal swelling  
transillumination

**Q2: What is the Dx?**

hydrocele

- Fluid filled sac ( fluid in a patent processus vaginalis or in the tunica vaginalis around the testicle).
- Communicating with the peritoneal cavity VS non communicating.
- **In most infants it will resolve in the 1<sup>st</sup> year.**
- If there is increase in size >> operation
- Any hydrocele appearing after a 1<sup>st</sup> year must be operated as it will not resolve.



# Undescended testicle

- Significant risks: infertility/ trauma/ torsion/ hernia/ cancer.
- Treatment : **orchidopexy** by the age of one year (6-12 months).
- After 2 years the testicle is abnormal and wouldn't be functioning.





## Q1: What is the Dx?


- Testicular torsion

## Q2: What is your Mx?

- Orchiectomy

### DDx for Acute scrotum:

1. Testicular torsion.
2. Torsion of testicular appendages.
3. Epididymorchitis.
4. Scrotal edema.
5. Complicated hernia.



# Imperforate anus

- Males > females.
- High lesion vs. low lesion.
- Meconium or air per urethra or vagina.
- One of the common findings that the anal opening anteriorly located.
- Treatment : resuscitation/ the low types managed by a one stage procedure in the neonatal period (**anoplasty**).
- Other types treated by colostomy in the neonatal period followed by a definitive procedure called **pull-through (posterior sagittal anorectoplasty)**.






Skin



**Mole (Melanocytic nevus): increased no., abnormal clusters, normal or increased production**





**Q1: Name the Dx?**

- Melanoma



**Q2: What is the most accurate prognostic factor?**

- The Depth

**Q3: Increased melanin production with normal number of cells is known to cause?**

- Freckles

**Q4: Mention 2 staging systems?**

1) Clark's level

2) Breslow's thickness



# seborrheic keratosis

- in the elderly " aka senile warts "
- special diagnostic feature : because they are patches of thick squamous epithelium they can be picked off if you try to pick the edges with a blunt forceps.
- when it peels off , it leaves a patch of pale-pink skin with slight bleeding.
- no other skin lesion behaves like this.
- doesn't need surgery. Completely benign.



- If a nevus undergoes changes in the pigmentation or in the shape or ulceration it indicates a melanoma.
- We differentiate the nevus from the vascular anomaly by its color.







## Hairy nevus

- It's premalignant and must be surgically removed.
- Congenital.
- Black or brown pigmented area with excess hair growth.

- In general, hair tuft or lipoma or hairy nevus located at the lower end of the back, it is associated with spina bifida.





**Q: a patient with pain and fever:**

**Q1: What is the Dx?**

- Cellulitis

**Q2: What is the micro-organism causing this?**

- Group A streptococci (GAS – mc!), Staph. Aureus



# Erysipelas

1. usually caused by streptococcus bacteria (beta hemolytic group A ).
2. Erysipelas is more superficial than cellulitis.
3. It's typically more RAISED and DEMARCATED.
4. The infection may occur on any part of the skin including the face, arms, fingers, legs and toes, BUT IT TENDS TO FAVOR THE EXTREMITIES.
5. Fat tissue is most susceptible to infection, and facial areas typically around the eyes, ears, and cheeks.



**Q: a patient post-splenectomy due to RTA:**  
**Q1: What is the micro-organism causing this?**  
- Meningococcus

**Q2: How can you prevent it?**  
MCV Vaccine

Vaccine should be 14 days BEFORE surgery , and in case of emergency surgery like this case it should be as soon as possible after surgery not 14 days after, others said in elective surgeries, it should be given 14 days before the operation But in emergent surgeries, it should be given at least 14 days post operatively.

**Post-  
Splenectomy:**  
We Give MCV,  
PCV, HiB



# Post Splenectomy Vaccination

- **Non-elective**

- Non-elective splenectomy patients should be vaccinated on or after postoperative day 14.
- Asplenic patients should be revaccinated at the appropriate time interval for each vaccine.

- **Elective**

- Elective splenectomy patients should be vaccinated at least 14 days prior to the operation.
- Asplenic or immunocompromised patients (with an intact, but nonfunctional spleen) should be vaccinated as soon as the diagnosis is made.
- Pediatric vaccination should be performed according to the recommended pediatric dosage and vaccine types with special consideration made for children less than 2 years of age.
- When adult vaccination is indicated, the following vaccinations should be administered:
  - ***Streptococcus pneumoniae***
    - Polyvalent pneumococcal vaccine (Pneumovax 23)
  - ***Haemophilus influenzae* type B**
    - *Haemophilus influenzae* b vaccine (HibTITER)
  - ***Neisseria meningitidis***
    - Age 16-55: Meningococcal (groups A, C, Y, W-135) polysaccharide diphtheria toxoid conjugate vaccine (Menactra)
    - Age >55: Meningococcal polysaccharide vaccine (Menomune-A/C/Y/W-135)

Vaccine	Dose	Route	Revaccination
Polyvalent pneumococcal	0.5 mL	SC*	Every 6 years
Quadravalent meningococcal/diphtheria conjugate	0.5 mL	IM upper deltoid	Every 3-5 years <sup>†</sup>
Quadravalent meningococcal polysaccharide	0.5 mL	SC*	Every 3-5 years
Haemophilus b conjugate	0.5 mL	IM*	None

\*Administered in the deltoid or lateral thigh region.

<sup>†</sup>Contact the manufacturer for the latest recommendations prior to revaccination.



# Non melanoma skin cancer

- The most common type of cancer.
  - Its mortality is low.
  - 75% BCC and 25% SCC.
- BCC is slow growing, locally destructive and rarely metastasize.
  - 80% are on head and neck.
- Melanin is a protective against tumor so blacks are less to have skin tumors.

**Q: Lesion on the face <1cm:**

**Q1: What is the Dx?**

- Basal cell carcinoma (BCC)

**Q2: What is the MCC?**

- Long exposure to sunlight



**Q3: Mention 2 ways of Mx?**

A) Non surgical:

(topical immunotherapy, intralesional interferon INJ, photodynamic)

B) Surgical (Excisional or destructive):

- Destructive: cautery, curettage, cryotherapy, CO laser ablation

- Excisional: Moh's micrographic surgery (MMS), Wide local excision

**Q4: What is the safety margin?**

- 4-10mm

**Q5: Write an alternative Mx?**

- Moh's micrographic surgery (MMS)



### Q6: Name 2 complications?

- METS, Ulceration

### Q7: Potential METS rate:


- <0.55 (from google)

### Q8: Do you expect to find enlarged LN?

- No (local disease)

### Q9: What does the arrow indicate?


Rodent ulcer (complication of BCC)



- Arising in the germinating basal cell layer of epithelial cells.
- **Nodular** (ulceration, telangiectasia, pearls).
- **Morphea** (manysites at the same time/ more aggressive than the nodular type).
- Slow growing.
- Local (rare risk of metastasis).

**Q: What is the type of cancer seen in this histology (biopsy taken from the nose tip):**

**- Basal Cell Carcinoma**





**Q: A 75 year old male farmer, heavy smoker presented with this lesion.**

**Q1: What is the most probable Dx?**

Squamous cell carcinoma.

**Q2: What is the LN of this area?**

Submental and submandibular??

**Q3: What will you do to confirm Dx?**


Biopsy for histopathology.



Basal cell carcinoma



Squamous cell carcinoma




- Arising from epidermal cells.
- Risk factors: sun exposure/ pale skin/ arsenic/ xeroderma pigmentosum/ immunosuppression.
- **Actinic keratosis : the precursor skin lesion.**
- Raised, slightly pigmented skin lesion/ ulceration/ exudate/ itching.
- Dx: excisional biopsy for small lesion/ incisional biopsy for large lesions.
- Most common sites : head, neck and hand.
- **Involves the lower lip and BCC involves the upper lip or above this level.**

**Q1: Name the lesion?**

- Onion cluster cells

**Q2: Mention the Dx?**

- SCC (Squamous cell carcinoma)





**Q: Two patients came to ER  
complaining of neck swelling:**

**Q1: What is the pathology?**

- Carbuncle

**Q2: MCC?**

- Staphylococcus Aureus

**Q3: Mx?**

- Incision, drainage and antibiotics



Carbuncle is an abscess larger than furuncle, usually with one or more openings draining pus onto the skin



**Q1: Identify this picture:**

Furuncle

**Q2: Mention one risk factor?**

DM

**Q3: it is more common in?**

In the back of the neck

**Q4: Name 1 treatment?**

Incision and drainage plus  
antibiotics







**actinic  
keratosis**



**Keratoacanthoma**  
self limiting growth and  
subsequent regression of  
hair follicle cells

**Q1: Dx of picture (1)?** Keratoacanthoma

**Q2: Dx of picture (2)?** Actinic Keratosis


**Q3: Dx of picture (3)?** Seborrheic Keratosis

**Q4: Dx of picture (4)?** Necrobiosis Lipodica

**Q5: Which doesn't have pre-malignant potency?**

3

**Q6: Picture 2 can convert to? SCC**




**Q1: What is this?**

- Lipoma

**Q2: What is the risk of wound infection after removal (% of wound infection)?**

- 1-3% (clean wound)





**Q: Give 2 DDx of a scalp lump?**

- 1) Sebaceous cyst
- 2) Epidermoid cyst






# Sebaceous cyst

- Benign subcutaneous cyst filled with sebum.

- found in hairy areas (scalp, scrotum, neck, ...).

- Most small cysts do not require treatment. Large or painful cysts may be removed surgically or by liposuction.

Important note: if there is a scalp lesion like this it's impossible to be lipoma as a differential diagnosis since lipoma emerges from fat under the skin and scalp area is devoid from fat.



Dr Vikram Singh



# Lipomatosis

AD condition in which multiple lipomas are present on the body.





**Q1: Describe what you see?**

- 1) Café au lait macules
- 2) Neurofibromas

**Q2: What is your Dx?**

- Neurofibromatosis

**Q3: Mention type of inheritance?**

- Autosomal Dominant





**Q: what is this and where do we find it??**

**A: Suppurative Hydradinitis** in axilla Found in sites of apocrine glands: axilla ,buttocks and perineum etc.

- caused by staph. Aureus.
- Treatment : antibiotics/ excision of skin with glands for chronic infection.



# Gas Gangrene

- Caused by *Clostridium perfringens*.
- Surgical emergency.



# Contusion

- Bruising injury caused by blunt trauma.
- Small hematoma is resorbed by itself (except on the face; need to be opened and evacuated)
- Large hematomas : if <24 hrs managed by aspiration, if > 24 hrs by incision and drainage.



# Abrasion

Managed by dressing to prevent 2ry bacterial infection.



**What is the type of this wound ? How is it treated?**

It's an **incised wound**.

Within the first 6 hours (or the first 24 hours in the face) it's treated by primary closure if the edges can be approximated without tension.



**Lacerated wound** usually caused by blunt objects.

First, we clean the edges (wound excision) to transform it to incised wound, then if within first 6 hours without contamination we close it by closure if the edges can be approximated without tension.

## Puncture wound

- Caused by pointed objects.
- Management: tetanus vaccine/ excision/ removal of foreign bodies.



---

## Avulsion flap

- Undermined laceration in the dermis and subcutaneous tissue.
- Management: debridement of edges/ excision of small avulsion flaps to prevent **trap-door effect**/ suturing.







## pyogenic granuloma

- During wound healing if the capillaries grow too vigorously they may form a mass covered with epithelium.
  - Look for a history of trauma
  - Very rapid growth

# Keloid Scar



# Hypertrophic Scar



Hypertrophic scar	Keloid scar
Improves with time (2 years)	No improvement with time
No genetic predisposition	Genetic predisposition
Less collagen	More collagen
Less cytokines	More cytokines
fibers parallel to the dermis	Fibers random in orientation
Remains within the borders of the original scar	Extends beyond the original scar margins
Regress spontaneously or by medication	

### Treatment :

- Surgery (Z- plasty, W- plasty) / artificial skin/ steroids/ pressure therapy/ topical silicon/ low dose radiation/ laser (CO2 and argon)/ calcium channel blockers/ interferon.

## Q1: Name the Dx?

- Keloid

## Q2: Name 2 RF?

- 1) Dark skin
- 2) FHx

## Q3: Name two characteristics?

- 1) Extend beyond borders of original wound
- 2) More common in darker skin
- 3) Require years to develop
- 4) Thick collagen







## **Granulation tissue**

(sign of healing ulcer)

# Inspection.....

## ○ Edge: five types:-

- *Sloping edge* e.g. healing ulcer
- *Punched out edge* e.g. Gummatous ulcer, deep trophic ulcer
- *Undermined edge* e.g. tuberculous ulcer-destroy subcutaneous faster the skin
- *Raised edge* e.g. Rodent ulcer
- *Rolled out (everted)*- e.g. Squamous Cell Carcinoma

Sloping  
(a healing ulcer)



Punched-out  
(syphilis, trophic)



Undermined  
(tuberculous)



Rolled  
(basal cell carcinoma)



Everted  
(squamous cell carcinoma)



**Figure 1.15** The varieties of ulcer edge.




## Q1: Name the Dx?

- DM/Peripheral arterial disease

## Q2: Causes?

- Prolonged pressure
- Uncontrolled long standing DM



### **Neurotrophic Ulcers:**

punched-out appearance  
painless.

Muscle atrophy may be noted.

**Q1: What is the most common etiology of this ulcer.**

- Neuropathic Diabetic Ulcer

**Q2: What is the most important step to accelerate healing?**

- Diabetic control, Decrease pressure at the area, Try to prevent infection and increase perfusion to the area





## Marjolin ulcer (malignant ulcer)

- SCC arises in a long standing benign ulcer or scar (long standing venous ulcer or scar of old burn ).
- Need 20-30 years to develop.



## Pressure sores grades

- 1) Erythema for >1 hour after relief of pressure ( Hyperemia).
- 2) Blisters with break in dermis, erythema requires 36 hr to disappear when relieved.  
( Ischemia, pressure 2-6h).
- 3) SC tissue and muscle involvement, skin is blue and thick ( Necrosis, pressure > 6 h).
- 4) Bone and tendon involvement, frank ulcer develops.



## Surgical treatment of pressure sores

- 1 excisional debridement.
- 2 partial or complete osteotomy.
- 3 closure of the wound with healthy, durable tissue. Closure can be either :
  - direct closure (in very small pressure sores).
  - skin grafts.
  - flaps.

### Flaps :

- Local tissue flaps.
- Myocutaneous flaps.
- Fasciocutaneous flaps.

**Contributing factors :** 1- pressure. 2- immobility. 3- shear (tangential pressure). 4- moisture. 5- malnutrition.



**Ischial pressure sore**



**Sacral pressure sore**



**Trochanteric pressure sore**

**Q: An 80 year old, bedridden male had this lesion in the buttock and lower back area.**

**Q1: What is this lesion?**

Pressure ulcer (bed sore)

**Q2: What is the most common cause?**

Pressure?





# Frost bite

- **Tissue freezing injury.**
- Mc type of cold injury.
- **At temperature (-2c).**
- **Treatment: rapid warming (40-42 C)/** debridement of clear blisters whereas hemorrhagic are left intact and aspirated if infected / elevation/ topical thromboxane inhibitor/ NSAID.
- Massage is contraindicated.



# Chilblains

- a type of **non-freezing tissue injury.**
- caused by chronic high humidity and low Temp with normal core Temp.
- seen commonly in mountain climbers.





# Trench foot

- The extremities are exposed to damp environment over long periods at temperatures ( 1- 10 C).
- Numbness/ tingling/ pain/ itching.
- The skin initially red and edematous then gradually turns to gray-blue discoloration.
- **Non- tissue freezing injury.**





**Pernio** is an inflammatory skin condition presenting after exposure to cold as pruritic and/or painful erythematous-to-violaceous acral lesions. Pernio may be idiopathic or secondary to an underlying disease.

- **Non tissue freezing injury.**



## Cold urticaria

- Familial and acquired.
- History of cold stimulation.



# Fight bite


\* over the dorsal metacarpophalangeal (MCP).

\* **organism** : *Eikenella corrodens* (specific to human mouth).

\***Complications**: cellulitis; extensor tenosynovitis; septic arthritis .

## \***Management**:

- 1) exploration (foreign body +extent)
- 2) local anesthesia
- 3) debridement
- 4)admission : drainage + ( IV) antibiotics (amoxicillin +clavulanic acid )



# Fournier Gangrene

necrotizing fasciitis in the perineum.

most commonly caused by c.perfringes.

Treat with tissue debridement and antibiotics.





# Kaposi sarcoma

- malignant proliferation
- associated with **HHV-8**.
- **Classically seen in three groups:**
  - 1) Transplant recipient, early spread, Rx decrease immunosuppression.
  - 2) older eastern European males, remain localized, Rx surgical removal.
  - 3) AIDS( Aids defining disease) - tumor spreads early, Rx increase antiretroviral therapy.



(cutaneous sarcoma appears as red hemispherical nodules or plaques)  
- is it painful ? no it is painless  
- usually associated with what ? HIV infection & AIDS



### **felon (whitlow):**

distal pulp space infection  
, if not treated results in  
osteomyelitis.  
cause : pricking.



### **Paronychia:**

infection of the nail fold ,  
happens due to bad manicure  
or bad maneuvering of  
hangnails.  
Most common hand infection.



## Tenosynovitis

- Infection of the synovial sheath surrounding tendon.

- The most causative organism of hand infection (tenosynovitis, felon, paronychia) is staph. Aureus.
- The 2<sup>nd</sup> is streptococcus.
- Initial treatment : oxacillin/ampicillin.
- Then we do culture and give antibiotics of choice.
- If abscess formed, incision and drainage.
- Elevation to decrease the edema.
- Resting the organ to decrease the pain.

# Antibioma

Hard, edematous swelling containing **sterile pus** following the treatment of an abscess with long term antibiotics rather than incision and drainage.

Treatment: exploration & drainage if it is indistinguishable from a carcinoma, otherwise spontaneous resolution takes place over several weeks.





# Bowen's disease



# Nevoid Basal Cell Syndrome

( AD )

Presentation :

- 1)multiple BCC mostly on the face
- 2)Cysts in the jaw.
- 3)Intracranial calcifications.
- 4)Rib abnormality ( mostly bifid ribs).



# Xeroderma pigmentosa

- It might predispose to SCC.
- an inherited premalignant condition associated with increase risk of all types of skin tumors.
- defect in the DNA repair genes
- AR





# Skin graft

**Q: What are the signs of graft take?**

1. The graft is adherent to the recipient site.
2. Pink color.
3. The graft blanches with pressure (denotes vascularity).





# Skin grafts

## 1- split thickness skin grafts :

- Epidermis and thin part of dermis.
- The donor site heals by epithelialization within 2 weeks.
- Used for large areas.




## 2- full thickness skin grafts:

- Taken from areas of loose skin as the donor area is closed by approximation of the edges (direct closure).
- Used for small areas.





- This is dermatome.
- It's used for taking a split thickness skin graft.



Split thickness skin graft after it has been meshed, showing the small perforations that allow the graft to be expanded and cover a greater area and also allows any blood/serum to drain away.

# Flaps

- A flap is a piece of tissue carries its own blood supplies that is moved from its original site, to cover a defect.
- Skin flaps/ muscle flaps/ myocutaneous flaps/ fasciocutaneous flaps/ osseofasciocutaneous flaps.
- Flaps are used when grafts are insufficient to cover the defect, or they wouldn't be taken.
- **To cover an avascular area.**
- When we need a more bulky tissue to deal with the defect and skin is not enough.
- **The donor area is managed by approximation if it was loose or by skin graft.**





# Burns

A dramatic night scene of a large fire consuming a building. The fire is intense, with bright orange and yellow flames reaching high into the dark sky. Thick black smoke billows from the burning structure. In the foreground, the silhouettes of four firefighters wearing helmets are visible, looking towards the inferno. The overall atmosphere is one of a major emergency response.



# 1st, 2nd, and 3rd Degree Burns





## 1<sup>st</sup> degree burn

- Pain and erythema.
- Limited to the dermis.
- No contracture.
- (1-6) days , heals by regeneration.
- Applies only to thermal burns.



## 2<sup>nd</sup> degree burn

- Necrosis of the epidermis and varying depth of the dermis (superficial/ intermediate/ deep).
- Pain, erythema, blisters, blanching, burned area is wet with exudate.
- Applies only to thermal burns.



## 3<sup>rd</sup> degree burn

- Full thickness.
- Eschar (dead tissue, insensitive, lethargy, inelastic, hard).
- Applies only to thermal burns.



- **Post burn contracture.**
- a complication of 3rd degree burns.
- they should have put skin graft for the patient to prevent this complication.




# Table 1. Classification of Burns by Depth

Burn Thickness	Deepest Skin Structure Involved	Appearance	Pain	Prognosis (Without Surgical Intervention)
Superficial (first-degree)	Epidermis	Dry, blanching erythema	Painful	Heals without scarring, 5-10 days
Superficial partial-thickness (second-degree)	Upper dermis	Blisters; wet, blanching erythema	Painful	Heals without scarring, < 3 weeks
Deep partial-thickness (second-degree)	Lower dermis	Yellow or white, dry, nonblanching	Decreased sensation	Heals in 3-8 weeks; likely to scar if healing > 3 weeks
Full-thickness (third-degree)	Subcutaneous structures	White or black/brown, nonblanching	Decreased sensation	Heals by contracture > 8 weeks; will scar

First degree	Partial thickness burns. <ul style="list-style-type: none"> <li>✱ Characterized by erythema (localized redness).</li> <li>✱ Appear sunburn-like.</li> <li>✱ Are not included when calculating burn size.</li> <li>✱ Usually heal by themselves.</li> </ul>	Third degree	Full thickness burns. <ul style="list-style-type: none"> <li>✱ Full skin has been destroyed.</li> <li>✱ Deep red tissue underlying blister.</li> <li>✱ Presence of bloody blister fluid.</li> <li>✱ Muscle and bone may be destroyed.</li> <li>✱ Require professional treatment.</li> </ul>
Second degree	Partial thickness burns. <ul style="list-style-type: none"> <li>✱ Part of skin has been damaged or destroyed.</li> <li>✱ Have blisters containing clear fluid.</li> <li>✱ Pink underlying tissue.</li> <li>✱ Often heal by themselves.</li> </ul>	Fourth degree	Full thickness burns. <ul style="list-style-type: none"> <li>✱ Penetrate deep tissue to fat, muscle, bone.</li> <li>✱ Require immediate professional treatment.</li> </ul>



# Role of 9's in Burns



# Parkland Formula

Volume of Lactated Ringers solution:

$$4\text{ml} \times \text{BSA}(\%) \times \text{weight}(\text{kg})$$

Give half of the  
solution for the

**first 8 hours**

Give the other half  
of the solution for the

**next 16 hours**

**Q: What is the Dx?**

- 2<sup>nd</sup> degree burn



**Q1: What is the degree of burn in this image?**

- 3<sup>rd</sup> Degree

**Q2: What is the name of the scar?**

- Escharotomy

**Q3: if the burn was circumferential and the patient weight was 100 kg, calculate:**


**1. TBSA%:**

- 100% (all the areas affected!)

**2. Fluid that needed in the 1<sup>st</sup> 8 hours if the TBSA is 40%:**

- 8 L


( $4 \times 40 \times 100 = 16\text{K ml}/1000 = 16 \text{ L}$ , in the 1<sup>st</sup> 8 hr we give  $\frac{1}{2}$  (so 8))





# Escharotomy VS fasciotomy

- fasciotomy is done in Mx of compartment syndrome after electrical burn.
- Escharotomy is done to decompress tissues in 3rd degree burns.
- Beneath escharotomy you will see granulation tissue, beneath fasciotomy you will see muscles.
- If ischemia is suspected, escharotomy is indicated.



# Electrical burn

- The severity depends on the voltage.
- Nerves, muscles and blood vessels have low resistance, so they are affected most.
- Skin, bone and tendons have high resistance, hence, they are less burned.
- **Management:**
  - ✓ Pt should be monitored for cardiac arrhythmias.
  - ✓ Good hydration & alkalization of urine to prevent renal impairment.
  - ✓ Fluid management couldn't be based on calculated formula.
  - ✓ Observation of limb vascularity & fasciotomy.

**What is the Dx?** Electrical burn

**What to do?** Fasciotomy.

**What is the cause of urine color?** Myoglobin.  
(electrical burn causes myoglobinuria)



# Thermal Burn

- Temperature  $> 45$  degrees.
- Duration of exposure is more important than degree of temp.
- Classification:
  - 1) direct flame burn
  - 2) scald burn (with hot liquids).
  - 3) contact burn with hot metals.
  - 4) friction burn.



**Scald burn**



**Contact burn**



**Friction burn**



**Q1: What category of burn does this patient have?**

-It's a facial flame burn ( facial edema ).

**Q2: What is the main risk of this burn?**

-the patient will have upper airway obstruction and risk of CO poisoning.

**Q3: What should you do?**

-The patient should be intubated before reaching to complete obstruction and give 100% oxygen if CO poisoning is suspected.





**Q: This lady had a flame burn 2 years ago.**

**Q1: What does the image show?** Post-burn fibrosis and contracture.

**Q2: What was the degree of her burn?** 3<sup>rd</sup> degree.

**Q3: Name the most suitable type of skin graft to use in reconstruction?**  
Full thickness

**Q: Serious complication that you fear from?** Transformation into SCC



**Q: This baby presented to the ER with scald burn.**

**Q1: What is the degree of burn?** 2<sup>nd</sup> degree.

**Q2: Mention three lines of acute Mx of the burn:**  
Fluid resuscitation/ pain control/ dressing.



# Chemical burns

- Caused by acids or alkali.
- **Acids** produce **less** damage and **less** penetration.
- **Acids** produce **coagulative** necrosis.
- **Alkali** produce **liquifactive** necrosis.
- Management : dilution by water for 2-4 hrs in alkaline burn, and 30 minutes for burns caused by acids.





A photograph of surgeons in an operating room, viewed from above. The surgeons are wearing blue scrubs, masks, and caps. They are holding surgical instruments, and a large surgical light is visible in the background. The text "General Surgery & Others" is overlaid in the center.

# General Surgery & Others




**Q: A trauma pt presented to the ER and was assisted with FAST:**

**Q1: What does FAST stand for?**

- **F**ocused **A**ssessment with **S**onography for **T**rauma

**Q2: What are the 4 sites that we look at in FAST?**

- 1) RUQ (Morison's pouch – Perihepatic)
- 2) LUQ (Perisplenic area)
- 3) Subcostal (Pericardiac)
- 4) Pelvic space



**Q: A patient presented to the ER  
after RTA:**

**Q1: What's your 1<sup>st</sup> priority?**

- ABC

(some said only airway)

**Q2: What's your 2<sup>nd</sup> priority?**

- Stop bleeding

(some said only breathing)



# Bleeding Classes

Parameter	Class I	Class II	Class III	Class IV
Blood loss (ml)	<750	750-1500	1500-2000	>2000
Blood loss (%)	<15	15-30	30-40	>40
Pulse rate (beats/min)	<100	100-120	>120	>140
BP	Normal	Decreased	Decreased	Decreased
Respiratory rate	14-20	20-30	30-40	>40
Urine output (ml/h)	>30	20-30	5-15	Negligible
CNS symptoms	Normal	Anxious	Confused	Lethargic

CNS: Central nervous system, BP: Blood pressure

**Q: This patient arrived to your ER after being stabbed as shown 15 minutes ago. He was anxious and his vital signs were BP: 95/55 mm Hg, pulse 105 BPM, and RR 25 Per minute.**

- 1. What is his class of hemorrhage? Stage 2**
- 2. How much blood has he lost? 750-1500 ml**





**Q: A patient fell and broke her leg, then the doctor who saw her put a cast on the leg, afterwards she complained from pain, swelling, redness and numbness in the same limb:**

**Q1: What is the Dx?**

- Compartment Syndrome

**Q2: Next step in Mx?**

- Decompression
- Remove the cast
- Fasciotomy

## **Q1: Name this sign?**

- Seat belt sign

## **Q2: Name 4 associated injuries?**

1) Flail chest

2) Small bowel injury

3) Cervical spine injury

4) Fracture of the sternum, ribs, clavicle & the vertebral bodies



**Q1: In penetrating trauma most affected organ?**


- Liver

**Q2: What type of injury more severe (blunt or penetrating)?**

- Blunt

**Q3: In a penetrating wound, what should you do?**

- Exploration Surgery



## *Blunt Vs Penetrating abd. Trauma...*

- Blunt trauma
  - spleen (45%)
  - liver (40%)
  - *Small bowel (10%)*
- Penetrating injuries
  - Stab wounds:-
    - the liver (40%),
    - small bowel (30%),
    - diaphragm (20%),
    - colon (15%);
  - gunshot wounds
    - small bowel (50%),
    - colon (40%),
    - liver (30%), and
    - vessels (25%).



# Abdominal injury- Evisceration



**Q: picture of multiple abdominal bruises, he asked about the zones of retroperitoneal bleeding and types of hemorrhage and where is the least likely place to check and when to go for surgery:**

- Traumatic retroperitoneal hematomas divided into 3 zones:

**Zone 1:** Centrally located, associated with pancreaticoduodenal injuries or major abdominal vascular injury

**Zone 2:** Flank or perinephric regions, associated with injuries to the genitourinary system or colon

**Zone 3:** Pelvic location, frequently associated with pelvic fractures or ileal-femoral vascular injury

- **Indication for exploration in retroperitoneal hematomas :**  
mandatory exploration should be performed in retroperitoneal hematomas resulted from penetrating injury, but the selection of treatment mode in blunt injury depend on the anatomical position of hematoma, visceral injury and the hemodynamic status of patients.

**Q: Hx of surgery for diverticulitis before 10, the amount collected over 24 hr is 1500 cc:**

**Q1: What is the pathology?**

- Enterocutaneous fistula  
(high output)

**Q2: What is the complication?**

- 1) Electrolyte disturbance
- 2) Skin excoriation
- 3) Sepsis

**Q3: What is the prognosis?**

- In most patients it closes spontaneously





**Q1: Type of stoma?**

- End Colostomy

**Q2: Mention 2 indications?**

- IBD

- Rectal Tumors







**Q: What is the complications in A, B, C?**

A) Prolapsed Stoma

B) Infected Stoma

C) Stoma Necrosis



**Q: A 65 year old man underwent abdomino-perineal resection 2 years ago after diagnosis of rectal ca.**




**Q1: What is the type of his stoma?**

End colostomy.

**Q2: What is the complication shown?**

Prolapse.





- Usually at the RLQ.
- Bag contents : watery stool.
- Offensive smell.
- Surrounding skin is usually inflamed (irritated from acid).
- Median or paramedian scar is usually seen.

**Loop ileostomy**  
2 openings

**End ileostomy**

## **Q1: What is this?**

Ileostomy.

## **Q2: How can you confirm?**

By its site and skin irritation around the stoma.

## **Q3: What is the disease that probably was treated by this?**

Chron's disease.

### **End Ileostomy**

- Edges are spouted.
- Site: right iliac fossa.








Loop colostomy

## End colostomy

- Sites : LLQ (sigmoid colon)/ RUQ (transverse colon) / RLQ (cecostomy)
- Formed stool in bag.
- No skin changes.
- Sigmoid colostomy expels stool 1/day.

Double barrel colostomy : together on left picture and separated on right picture.


## Double-barreled stoma





## incisional hernia

(notice the surgical scar)  
m.c.c is wound infection



## Femoral hernia

- most common hernia in females.
- Medial to femoral vessels.

## Q1: Name of the test?


- Ring occlusion test

## Q2: If you ask the patient to cough while you maintain pressure and you notice a bulge, what is your Dx?

- Direct inguinal hernia

\*\* Note: Ring occlusion test differs from 3 fingers test, You Ask the patient to cough> Impulse felt on the index finger> Indirect hernia So; Zieman's Test (3 Finger Test) is used to differentiate type of hernia.

- Index: deep inguinal hernia (indirect)
- Middle: superficial inguinal (direct)
- Ring: Saphenous opening (femoral hernia)



Indirect Inguinal Hernia	Direct Inguinal Hernia
Pass through inguinal canal.	Bulge from the posterior wall of the inguinal canal
Can descend into the scrotum.	Cannot descent into the scrotum.
Lateral to inferior epigastric vessels.	Medial to inferior epigastric vessels.
Reduced: upward, then laterally and backward.	Reduced: upward, then straight backward.
Controlled: after reduction by pressure over the internal (deep) inguinal ring.	Not controlled: after reduction by pressure over the internal (deep) inguinal ring.
The defect is not palpable (it is behind the fibers of the external oblique muscle).	The defect may be felt in the abdominal wall above the pubic tubercle.
After reduction: the bulge appears in the middle of inguinal region and then flows medially before turning down to the scrotum.	After reduction: the bulge reappears exactly where it was before.
Common in children and young adults.	Common in old age.



# Inguinal hernia

## DDx of inguinal hernia :

Hydrocele/ saphena varix/  
testicular torsion/ psoas  
abscess .. Etc.

- **Indirect** : most common type in both males and females.
- **Indirect** : lateral to the inferior epigastric artery.
- **Direct** : medial within hesselbach's triangle.



**Herniotomy** : only in peds patients.

**Herniorrhaphy** : tension due to approximation/  
high recurrence.

**Hernioplasty** : using a mesh/tension free/ open  
or laparoscopic.

# Para umbilical hernias

crescent-shaped bulge develops in the navel.




**Q: Patient presented with painful lump in his belly button:**

**Q1: What is the Dx?**

- Strangulated Hernia

**Q2: If the bowel still the same despite of all measures, what's your next step?**


- Resection and Anastomosis






Poland syndrome

Pectus excavatum  
( funnel chest )



## Ganglion cyst

- is a non-neoplastic soft tissue lump.
- It's **painful**.
- recurrence may occur after surgery.





# Lower extremity amputations

Indications : irreversible tissue ischemia & necrotic tissue/ severe infection / severe pain with no bypassable vessels, or if pt is not interested in a bypass procedure.



Bellow knee amputation



Above knee amputation



Syme's amputation

Through the articulation of the ankle with removal of the malleoli.



Transmetatarsal amputation



Ray amputation


Removal of toe & head of Metatarsal.

# Flail chest

Segment occurs when three or more contiguous ribs are fractured in two or more places.

It typically occurs after high impact **trauma**.

Flail segment of chest wall that moves paradoxically (opposite to the rest of chest wall)



# DOG BITE

\*Management :

- 1)exploration
- 2)analgesia
- 3)IV antibiotics  
(clindamycin + penicillin)
- 4)elevation
- 5)tetanus toxoid
- 6)rabies vaccine





## Erythroplakia

- Reddish patch that appears on the oral mucosa.
- It has 17 X more risk of malignancy than leukoplakia.



## Leukoplakia

- White patch that appears on the oral or genital mucosa.
- Risk factors : smoking/ شرب
- Premalignant (transform to SCC).



## Q1: What is the Dx?

- Cushing Syndrome

## Q2: Causes?

- Iatrogenic (cortisol administration)
- Pituitary Adenoma

\*\* Note: Cushing triad:

- 1) Irregular, decreased respirations
- 2) Bradycardia
- 3) Systolic hypertension



**Q1: White arrow?**



- Pituitary Adenoma

**Q2: Syndrome name?**

- MEN

**Q3: The most important thing surgically to do for this patient?**

- Pancreatic tumor “not sure”



3P

2P 1M

1P 2M

## MEN 1

Pituitary  
adenoma

Parathyroid  
hyperplasia

Pancreatic  
tumors

## MEN 2A

Parathyroid  
hyperplasia

Medullary  
thyroid  
carcinoma

Pheo-  
chromo-  
cytoma

## MEN 2B

Mucosal  
neuromas

Marfanoid  
body  
habitus

Medullary  
thyroid  
carcinoma

Pheo-  
chromo-  
cytoma

- **MEN I (3 Ps)**


- Pituitary,
- Parathyroid,
- Pancreatic

- **MEN 2A (1M,2Ps)**

- MTC
- Pheochromocytoma
- Parathyroid

- **MEN 2B (2Ms,1P)**

- MTC,
- Marfanoid  
habitus/Mucosal  
neuroma
- Pheochromocytoma



**Q: Male with heart disease:**

**Q1: what is the abnormality in the picture?**

- Gynecomastia

**Q2: What drugs is the patient taking that might cause this?**

- Spironolactone

- Digoxin



## DRUGS CAUSING GYNECOMASTIA

Mnemonic: '**DISCKO**'


- **D**igoxin
- **I**soniazid
- **S**piro<sup>n</sup>olactone
- **C**imetidine
- **K**etoconazole
- **O**estrogen





# Charcot foot

- Rocker-bottom appearance.
- Develops as a result of neuropathy such as in diabetic pts.
- ttt : immobilization/  
custom shoes & bracing.



# signs of basilar skull fracture



Clear rhinorrhea



raccoon eyes




otorrhea



battle's sign ( ecchymosis  
behind the ear )



hemotympanum



**Hyphema:** blood in the anterior chamber of the eye

**Q: This is pelvic x-ray of a patient post RTA:**

**Q1: What is the pathology?**

- Pelvic fracture

**Q2: What is the most serious complication?**

- Bleeding (Femoral artery)





**Question: about post-operative fever:**

1. Lung Atelectasis
2. ECG change MI
3. UTI
4. wound surgical site infection
5. drugs

**Question A: which of the following picture are consider as a source of fever after 1-3 days?**


-Atelectasis (1)

**Question B: which of the following picture are consider as a source of fever after 5-7 days?**

-Wound infection (4)



**1**



**2**



**3**



**4**




**5**

Category	Day	Description
Wind	POD 1-2	the lungs, i.e. <a href="#">pneumonia</a> , aspiration, and pulmonary embolism; <a href="#">atelectasis</a> has been commonly cited as a cause of post-operative fever, but supporting evidence is lacking <sup>[2][3]</sup>
Water	POD 3-5	<a href="#">urinary tract infection</a> , possibly catheter-associated (if a urinary catheter was inserted during surgery or remains in place currently i.e. <a href="#">Foley catheter</a> )
Wound	POD 5-7	infection of the surgical incision(s), either superficial or deep <sup>[4]</sup>
(W)abscess	POD 5-7	infection of an organ or space <sup>[5]</sup>
Walking (or VEINS pronounced like "Weins")	POD 5+ (risk may persist for months post-operatively)	<a href="#">deep vein thrombosis</a> or <a href="#">pulmonary embolism</a>
Wonder drugs or "What did we do?"	Anytime	<a href="#">drug fever</a> or reaction to blood products, either a <a href="#">febrile non-hemolytic transfusion reaction</a> or <a href="#">transfusion-related acute lung injury</a>
Wing/Waterway	Anytime	bloodstream infection, phlebitis, or cellulitis related to intravenous lines, either central or peripheral

# Pre-cancerous lesions

1. Leukoplakia of the tongue (15 % malignant transformation to SCC / DDx: Oral candidiasis, how to differentiate? Candidiasis scrapes off).
2. Colon in FAP.
3. Colon in HNPCC.
4. Thyroid gland in MENS II.
5. Breast in BRCA mutations.

- Surgery has a role in 1ry cancer prevention.



# Classic physical findings that represent METS & incurable disease :

1) **Virchows node** enlargement (left supraclavicular nodes).



2) **sister mary josephs nodules** : infiltration of the umbilicus.




3) **blumers shelf** :fullness in the pelvic ,cul-de-sac(solid peritoneal deposit anterior to the rectum forming a shelf palpated on PR).

4) **krukenburgs tumor** :enlarged ovaries on pelvic examination (Metz to ovaries).

5) **hepatosplenomegaly** with ascites and jaundice.

6) **cachexia**.

7) **irishs node** :left axillary adenopathy.



Virchow's node enlargement



## Diagram of Tumour Markers

### Oesophagus

(CEA, SCC)

### Lung

parvicellular: NSE (CYFRA 21-1)  
non-parvicellular: (CEA, CYFRA 21-1)

### Liver/Biliary ducts

AFP, CA 19-9

### Bladder

(CYFRA 21-1)

### Uterus

SCC (CEA)

### Prostate gland

PSA

### Testes

AFP, HCG

### Thyroid gland

Thyroglobulin,  
Calcitonin (C-cell,  
CEA)

### Mamma

CA 15-3, CEA

### Stomach

CA 72-4 (CEA)

### Pancreas

CA 19-9 (CEA)

### Colorectal


CEA (CA 19-9)

### Ovaries

CA 125 (CA 72-4)

### Multiple Myeloma

$\beta_2$ -Microglobulin



A close-up photograph of a surgical team's hands in white gloves, positioned over a blue sterile drape. Various surgical instruments, including forceps and scissors, are laid out on the drape. The background is blurred, showing more of the surgical team in blue scrubs. The text "Tools & Instruments" is overlaid in a large, white, bold font with a black outline.

# Tools & Instruments

**Q1: What are the names of those tools?**

- Central line and cannula

**Q2: What is better to insert in a trauma patient & for fluid administration, why?**

- Cannula, because it is easier to use, require less experience and time, it also deliver the largest volume of fluid

**Q3: The smallest cannula in diameter is?**


- Purple

(Cannula's in the picture – Blue)


**Q4: Cannula for large amount of fluid?**

- Orange

(cannula's in the picture - Green)



**Triple Lumen**




**Double Lumen**



## IV NEEDLE GAUGES SIZE CHART


**14 GAUGE**



Color: Orange  
Outer Diameter: 2.1 mm

Length: 45 mm  
Flow Rate: 240 mL/min


**16 GAUGE**



Color: Gray  
Outer Diameter: 1.8 mm

Length: 45 mm  
Flow Rate: 180 mL/min


**18 GAUGE**



Color: Green  
Outer Diameter: 1.3 mm

Length: 32 / 45 mm  
Flow Rate: 90 mL/min


**20 GAUGE**



Color: Pink  
Outer Diameter: 1.1 mm

Length: 32 mm  
Flow Rate: 60 mL/min


**22 GAUGE**



Color: Blue  
Outer Diameter: 0.9 mm

Length: 25 mm  
Flow Rate: 36 mL/min


**24 GAUGE**



Color: Yellow  
Outer Diameter: 0.7 mm

Length: 19 mm  
Flow Rate: 20 mL/min

**26 GAUGE**



Color: Purple  
Outer Diameter: 0.6 mm

Length: 19 mm  
Flow Rate: 13 mL/min




**Q1: Name this tube?**

- Chest tube

**Q2: Give 4 indications?**


- 1) Hemothorax
- 2) Pneumothorax
- 3) Chylothorax
- 4) Empyema
- 5) Hydrothorax
- 6) Pleural Effusion
- 7) Post-op





Chest tube drain

Chest drain system



## **Q1: What is this device?**

- Nasogastric tube

## **Q2: Give 3 indications?**

- 1) Feeding
- 2) Decompression
- 3) Administration of medication
- 4) Bowel irrigation

## **Q3: The tip of it should reach?**

- Stomach body





## Q1: What is this?

- Gastric tube/G-tube/PEG tube/ Gastrestomy

## Q2: What is the main indication for it?

- Feeding






## Q1: What is A,B?

**A** > Stoma base (Flange)

**B** > Stoma bag

## Q2: Mention 3 indications?

- After proctocolectomy
- Imperforated anus
- Secondary healing
- Some said (colectomy, ileostomy, double barrel)



## **Q1: What is this?**

- Tracheostomy

## **Q2: Mention 2 complications?**

- 1) Infection
- 2) Blockage (Obstruction)
- 3) Bleeding
- 4) Pneumothorax

## **Q3: Mention 2 indications?**

- 1) Upper airway obstruction
- 2) Obtaining an airway in severe facial or neck trauma
- 3) Upper airway edema and copious secretions
- 4) Failure to wean from mechanical ventilation
- 5) Acute respiratory failure with need for prolonged mechanical ventilation  
(mc indication, 2/3 of all cases)




**Q1: Name the tube?**

- Nephrostomy tube

**Q2: Write 2 indications?**

- 1) Urinary obstruction secondary to calculi
- 2) Hemorrhagic cystitis






**Q1: Which one is not used in primary survey?**

- C (Foley's Catheter)

**Q2: Which one is your 1<sup>st</sup> priority?**

- D (Neck collar), some said (B)





**Q1: What is the name of device?**

- Foley's Catheter

**Q2: What is the unit used in measurement??**

- French



**Q1: What is this?** Colonoscopy

**Q2: Name 2 pathologic finding?**

- 1) Angiodysplasia
- 2) Diverticulosis
- 3) Colon tumor
- 4) Polyps, masses

**Q3: Name 2 therapeutic procedures done with it?**

- 1) Laser Ablation
- 2) Polyps Resection





**Q1: What is this device?**

- Pulse Oxymeter

**Q2: What does it calculate?**

- O2 Saturation
- Pulse Rate (HR)

**Q1: What is the name of the drain?**

- Penrose

**Q2: Type of the drain?**


- Open drain






# Q: Name of the drain?

- Corrugated Drain



# T-tube

used for post operative drainage of  
common bile duct.



# Redivac drain

Drains can be:  
**Open or closed**  
**Active or passive:**





### **Q1: What is this device?**

Intermittent pneumatic compression technique  
(Inflatable leg sleeves).

### **Q2: Uses?**

To prevent DVT.



**Q1: what is this?**  
incentive spirometer

**Q2: Why do we use it?**  
used after surgery to prevent atelectasis .  
(used while inspiration not expiration).





## Q1: Name of device seen in the CT?

- Inferior vena cava filter

## Q2: Give 1 indication for it?

- 1) Proven VTE with contraindication for anticoagulation.
- 2) Proven VTE with complications of anticoagulation.
- 3) Recurrent VTE despite adequate anticoagulation.



### **Q1: Name of device?**

- Central venous catheter (CVC)

### **Q2: Where do you insert it?**


- Subclavian vein
- Internal jugular vein

### **Q3: Mention 2 indications?**

- 1) Total parenteral nutrition (TPN)
- 2) Hemodialysis
- 3) Chemotherapy

### **Q4: Mention 2 complications?**

Pneumothorax, Hemothorax, Recurrent laryngeal nerve injury, Arterial or Venous injury, Arterial access instead of venous, Hematoma, Infection, Thrombosis and occlusion of the line...etc



# Venous access catheter

- Small, flexible hollow tube.
- Surgically placed into a large vein.
- Can be left for several months.
- Used for repeated infusions of **chemotherapy** drugs.




Venous access catheter

# Biological heart valves


Used in the following cases:

- Age > 60
- Previous thrombosed mechanical valve.
- Limited life expectancy.
- If Coagulation is contraindicated.
- Young women wishing to get pregnant.



# Mechanical prosthetic valves

Used if the age is < 60 + long life expectancy.






**Q: what can you see in  
this chest X-Ray ?**

sternal wires in the  
midline (indicate that  
patient U/W  
sternotomy).


pacemaker.



**Intra-aortic balloon pump (IABP)** is a mechanical device that increases myocardial oxygen perfusion and increasing CO. These actions combine to decrease myocardial oxygen demand and increase myocardial oxygen supply.


**Notes :**

- the polyethylene balloon has a radiopaque tip.
- the balloon inflates during diastole and deflates during systole .
- indications :Cardiogenic shock post-MI , (CABG) ,post cardiothoracic surgery, unstable angina .
  - most important complication is lower limb ischemia, we have to check the pulse and perfusion .
- most important contraindication: aortic valve insufficiency (AR) ,aneurysm .



A background image showing surgeons in blue scrubs and masks performing an operation. The scene is dimly lit with a bright light source focused on the surgical site.

Best of Luck!



دعواتكم